Product Datasheet

Integrin alpha V beta 5 Antibody (P5H9) [PE/Cy7] FAB2528PECY7

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/FAB2528PECY7

Updated 8/10/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/FAB2528PECY7



FAB2528PECY7

Integrin alpha V beta 5 Antibody (P5H9) [PF/Cv7]

Integrin alpha V beta 5 Antibody (P5H9) [PE/Cy/]	
Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark. Do not freeze.
Clonality	Monoclonal
Clone	P5H9
Preservative	0.05% Sodium Azide
Isotype	lgG1
Conjugate	PE/Cy7
Purity	Protein A or G purified from hybridoma culture supernatant
Buffer	PBS
Product Description	
Host	Mouse
Gene ID	3685
Gene Symbol	ITGAV
Specificity/Sensitivity	Detects human Integrin $\alpha V\beta 5$. Recognizes the human Integrin $\alpha V\beta 5$ heterodimer and does not recognize the αV subunit in association with any other β subunits.
Immunogen	HT1080 human fibrosarcoma cell line
Product Application Details	
Recommended Dilutions	Flow Cytometry
Application Notes	Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact

Const. 20. Const. (Const. Const. Cons
otimal dilution of this antibody should be experimentally determined. For stimal results using our Tandem dyes, please avoid prolonged exposure to light extreme temperature fluctuations. These can lead to irreversible degradation decoupling. When staining intracellular targets, specific attention to the fixation d permeabilization steps in your flow protocol may be required. Please contact ir technical support team at technical@novusbio.com if you have any testions.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/FAB2528PECY7

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

