# **Product Datasheet**

## Muscarinic Acetylcholine Receptor M3/CHRM3 Antibody (580011) [Allophycocyanin/Cy7] FAB6378APCCY7

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/FAB6378APCCY7

Updated 8/10/2024 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/FAB6378APCCY7

### FAB6378APCCY7

Muscarinic Acetylcholine Receptor M3/CHRM3 Antibody (580011) [Allophycocyanin/Cy7]

| Product Information          |  |
|------------------------------|--|
| Unit Size                    | 0.1 ml   |
| Concentration                | Please see the vial label for concentration. If unlisted please contact technical services.  |
| Storage                      | Store at 4C in the dark. Do not freeze.  |
| Clonality                    | Monoclonal   |
| Clone                        | 580011   |
| Preservative                 | 0.05% Sodium Azide   |
| Isotype                      | lgG2b  |
| Conjugate                    | Allophycocyanin/Cy7  |
| Purity                       | Protein A or G purified from hybridoma culture supernatant   |
| Buffer                       | PBS  |
| Product Description          |  |
| Host                         | Mouse  |
| Gene ID                      | 1131   |
| Gene Symbol                  | CHRM3  |
| Specificity/Sensitivity      | Detects HEK293 cells transfected with human CHRM3 in Cell-based ELISA  |
| Immunogen                    | NS0 mouse myeloma cell line transfected with human CHRM3<br>Accession # P20309   |
| Product Application Details  |  |
| <b>Recommended Dilutions</b> | Flow Cytometry   |
| Application Notes            | Optimal dilution of this antibody should be experimentally determined. For<br>optimal results using our Tandem dyes, please avoid prolonged exposure to light<br>or extreme temperature fluctuations. These can lead to irreversible degradation<br>or decoupling. When staining intracellular targets, specific attention to the fixation<br>and permeabilization steps in your flow protocol may be required. Please contact<br>our technical support team at technical@novusbio.com if you have any<br>questions. |

www.novusbio.com





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/FAB6378APCCY7

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

