## **Product Datasheet**

# SIRP alpha/CD172a Antibody (772734) [PE/Cy5.5] FAB7307PECY55

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/FAB7307PECY55

Updated 8/10/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/FAB7307PECY55



## **FAB7307PECY55**

SIRP alpha/CD172a Antibody (772734) [PE/Cy5.5]

| SIRP alpha/CD172a Antibody (772734) [PE/Cy5.5] |   |
|--|---|
| Product Information                            |   |
| Unit Size                                      | 0.1 ml  |
| Concentration                                  | Please see the vial label for concentration. If unlisted please contact technical services.   |
| Storage  | Store at 4C in the dark. Do not freeze.   |
| Clonality                                      | Monoclonal  |
| Clone  | 772734  |
| Preservative                                   | 0.05% Sodium Azide  |
| Isotype  | lgG2b   |
| Conjugate                                      | PE/Cy5.5  |
| Purity   | Protein A or G purified from hybridoma culture supernatant  |
| Buffer   | PBS   |
| Product Description                            |   |
| Host   | Mouse   |
| Gene ID  | 140885  |
| Gene Symbol                                    | SIRPA   |
| Specificity/Sensitivity                        | Detects rat SIRP-α/CD172a in direct ELISA and Western Blot.   |
| Immunogen                                      | Mouse myeloma cell line NS0-derived recombinant rat SIRP-α/CD172a Lys32-Asn373 Accession # P97710   |
| Product Application Details                    |   |
| Recommended Dilutions                          | Flow Cytometry  |
| Application Notes                              | Optimal dilution of this antibody should be experimentally determined. For optimal results using our Tandem dyes, please avoid prolonged exposure to light or extreme temperature fluctuations. These can lead to irreversible degradation or decoupling. When staining intracellular targets, specific attention to the fixation and permeabilization steps in your flow protocol may be required. Please contact our technical support team at technical@novusbio.com if you have any |



questions.



## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

# Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/FAB7307PECY55

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

