

Product Datasheet

Wnt-4 Antibody NB100-1006

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-1006

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-1006




NB100-1006**Wnt-4 Antibody**

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA
Product Description	
Host	Goat
Gene ID	54361
Gene Symbol	WNT4
Species	Human
Reactivity Notes	Predicted cross-reactivity based on sequence identity: Mouse, Rat.
Immunogen	Peptide with sequence C-SNWLYLAKLSSVGS corresponding to internal region according to NP_110388.2.
Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Peptide ELISA
Recommended Dilutions	Western Blot 1:100-1:2000, Immunocytochemistry/ Immunofluorescence 10 ug/ml, Peptide ELISA Detection limit 1:4000
Application Notes	WB: Approx. 40+35 kDa band observed in Jurkat lysates (calculated MW of 39.1 kDa band according to NP_110388.2 Negative Control: U937 cell lysate.). Immunofluorescence: Strong expression of the protein seen in the Vesicle of HeLa and MCF7 cells.




Images

Western Blot: Wnt-4 Antibody [NB100-1006] - (1ug/ml) staining of Jurkat (A) and negative control U937 (B) cell lysate (35ug protein in RIPA buffer). Detected by chemiluminescence.



Immunocytochemistry/Immunofluorescence: Wnt-4 Antibody [NB100-1006] - analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml), showing vesicle staining. The nuclear stain is DAPI (blue)



Publications

Huguet EL, McMahon JA, McMahon AP et al. Differential expression of human Wnt genes 2, 3, 4, and 7B in human breast cell. Cancer Res 1994-05-15 [PMID: 8168088]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-1006

NB820-59177	Human Brain Whole Tissue Lysate (Adult Whole Normal)
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1006

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



