

Product Datasheet

RKIP/PBP Antibody NB100-1020

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-1020

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-1020



NB100-1020

RKIP/PBP Antibody

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA

Product Description	
Host	Goat
Gene ID	5037
Gene Symbol	PEBP1
Species	Human
Reactivity Notes	Predicted cross-reactivity based on sequence similarity: Mouse, Rat, Canine, Bovine.
Immunogen	Peptide with sequence C-DYVPKLYEQLSGK corresponding to C-Terminus according to NP_002558.1.

Product Application Details	
Applications	Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA
Recommended Dilutions	Western Blot 0.5 - 2 ug/mL, Immunohistochemistry 5 ug/mL, Immunohistochemistry-Paraffin 5 ug/mL, Peptide ELISA Detection limit 1:32000




Images


Western Blot: RKIP/PBP Antibody [NB100-1020] - (1ug/ml) staining of A549 lysate (35ug protein in RIPA buffer). Detected by chemiluminescence.



Immunohistochemistry-Paraffin: RKIP/PBP Antibody [NB100-1020] - Negative Control showing staining of paraffin embedded Human Cortex, with no primary antibody.



Immunohistochemistry-Paraffin: RKIP/PBP Antibody [NB100-1020] - Staining of paraffin embedded Human Cortex. Antibody at 5 ug/mL. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



Publications

Hleihel R, Skayneh H, de The H et al. Primary cells from patients with adult T cell leukemia/lymphoma depend on HTLV-1 Tax expression for NF- κ B activation and survival Research Square 2023-03-08 [PMID: 37137914] (Flow Cytometry, Human)

Yeung K, Seitz T, Li S et al. Suppression of Raf-1 kinase activity and MAP kinase signalling by RKIP. Nature 1999-09-09 [PMID: 10490027]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-1020

NBL1-14274	RKIP/PBP Overexpression Lysate
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1020

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



