

Product Datasheet

HIF-1 alpha Antibody (H1alpha67) NB100-123

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 9 Publications: 161

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-123

Updated 10/23/2024 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-123




NB100-123

HIF-1 alpha Antibody (H1alpha67)


Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	H1alpha67
Preservative	0.05% Sodium Azide
Isotype	IgG2b
Purity	Protein A purified
Buffer	PBS with 1% BSA
Target Molecular Weight	93 kDa
Product Description	
Host	Mouse
Gene ID	3091
Gene Symbol	HIF1A
Species	Human, Mouse, Rat, Porcine, Avian, Bovine, Canine, Ferret, Primate, Rabbit, Sheep
Reactivity Notes	Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information. Rabbit reactivity reported in scientific literature (PMID: 16738327, 26339038).
Immunogen	This HIF-1 alpha Antibody (H1alpha67) was developed against a fusion protein containing amino acids 432 - 528 of human HIF-1 alpha [Uniprot# Q16665].
Product Application Details	
Applications	Western Blot, Chromatin Immunoprecipitation, ELISA, Flow Cytometry, Gel Super Shift Assays, Immunoblotting, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Chromatin Immunoprecipitation (ChIP), Knockdown Validated, Knockout Validated
Recommended Dilutions	Western Blot 1:500 - 1:1000, Chromatin Immunoprecipitation 1:10 - 1:500. Use reported in scientific literature, Flow Cytometry 1:10 - 1:1000, ELISA 1:100 - 1:2000. Use reported in scientific literature, Immunohistochemistry 1:100 - 1:300, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation 1:10, Immunohistochemistry-Paraffin 1:100 - 1:300, Immunoblotting reported in multiple pieces of scientific literature, Gel Super Shift Assays 1:1 - 1:100. Use reported in scientific literature, Chromatin Immunoprecipitation (ChIP) 1:10-1:500, Knockout Validated reported in scientific literature (PMID 27991597), Knockdown Validated
Application Notes	By WB, this antibody recognizes bands at 120kDa representing HIF-1 alpha in induced tissues and cells. Multiple bands may be seen at 120kDa representing post-translational modifications. Nuclear extracts are recommended for WB.

Images


Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Results of in situ hybridization and immunohistochemistry on thin adjacent section to detect expression of HIF-1a1.2 mRNA and HIF1a protein in malignant and benign prostate tissue. In situ hybridization (antisense probe, Fig. 3a) and immunostaining with HIF1a Ab2 (Fig. 3c) on thin adjacent sections of NE-differentiated prostate adenocarcinoma showed co-localization of HIF1a1.2 transcript and HIF-1a protein. Incubation with sense probe did not generate any detectable hybridization signals (Fig. 3b). Both In situ hybridization (Fig. 3d antisense) and HIF-1a Ab2 immunostaining (Fig. 3f) were negative in non-NE-differentiated prostate adenocarcinoma. In situ hybridization with sense probe performed on non-NE-differentiated prostate cancer (Fig. 3e). Image collected and cropped by CiteAb from the following publication (<https://www.biomedcentral.com/1471-2407/10/385>), licensed under a CC-BY license.




Knockdown Validated: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - HIF-1 inhibits the apoptosis of hypoxic glioblastoma cells. U87MG cells were transfected with siRNA against HIF-1alpha. Forty-eight hours after transfection, cells were incubated for 2 h in hypoxia (1% O₂). HIF-1alpha was detected by immunoblotting. beta-Actin was used as a loading control. The results are representative for three independent experiments. Image collected and cropped by CiteAb from the following publication (<https://www.nature.com/articles/cddis2013562>) licensed under a CC-BY license.




Western Blot: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - MLN4924 induces accumulation of HIF1a in a time dependent manner. Cells were treated with 0.1uM MLN4924 for indicated time periods, followed by IB with indicated antibodies. Image collected and cropped by CiteAb from the following publication (<https://www.nature.com/articles/cddis2012125>) licensed under a CC-BY license.




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Analysis of HIF-1 alpha in human lung tissue. Image courtesy of product review by Aneta Gandjeva.



Immunoprecipitation: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - MG132 attenuates the inhibitory effect of interleukin-1beta on HIF-1/AM axis. U87MG cells were incubated for 2 h in hypoxia (1% O₂) with or without interleukin-1beta (10 ng/ml). The proteasome inhibitor MG132 was added 30 min in advance to prevent ubiquitinated HIF-1alpha from proteasomal degradation. HIF-1alpha was precipitated by anti-HIF-1alpha antibody. Ubiquitin was detected by immunoblotting. Non-specific mouse IgG was used as a negative control. The results are representative for three independent experiments Image collected and cropped by CiteAb from the following publication (<https://www.nature.com/articles/cddis2013562>) licensed under a CC-BY license.




Western Blot: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Analysis using the HRP conjugate of NB100-123. Detection of HIF-1 alpha in cobalt chloride treated/untreated COS-7 nuclear extracts using NB100-123.




Western Blot: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Analysis using the HRP conjugate of NB100-123. On day 1, MEF cells (+/+, -/-), were grown on 15cm dish (2x10 to the 6th cells). On day 2, cells were exposed to hypoxia for 4hrs. Cells were washed with ice cold PBS twice and whole cell protein was extracted extracted with RIPA buffer fortified with protease. Upon quantification, 100ug of protein was fractionated on 7% polyacrylamide gel. Gel was transferred overnight onto nitrocellulose membrane. The membrane was probed with HIF-1 alpha monoclonal antibody at a 1:500 dilution (NB100-123). The secondary antibody was conjugated with HRP and was used at a 1:2500 dilution. Photo courtesy of Dr. Gregg Semenza, Johns Hopkins University.




Immunocytochemistry/Immunofluorescence: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Dapagliflozin protects against metabolic switch by inhibition of HIF-1 alpha. Representative images of PTCs incubated with 10 umol/l of molidustat for 6 h with immunofluorescent staining for HIF-1 alpha. Scale bar = 10 um. Image collected and cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/33187129/>) licensed under a CC-BY license.



Immunohistochemistry-Paraffin: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunohistochemical detection of HIF-1alpha protein in human tissues. The nature of the tissue is indicated on top of each figure. Original magnifications are as follows: A, x400; B, x400; C, x100; D, x100; E, x100; F, x100; G, x40; H, x40. Image collected and cropped by CiteAb from the following publication (<https://bmccgenet.biomedcentral.com/articles/10.1186/1471-2156-5-27>), licensed under a CC-BY license.




Immunocytochemistry/Immunofluorescence: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Staining in pig endothelial cells under hypoxia condition using NB100-123. Image from verified customer review.




Immunohistochemistry-Paraffin: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Staining on pig tissue (brown) using NB100-123. Image from verified customer review.




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Staining of biotin conjugated HIF-1 alpha (NB100-123B) in human kidney.




Immunoprecipitation: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunoprecipitation of HIF-1alpha using NB100-123 in COS7 CoCl₂ treated lysate. Heavy and light chains are also detected. Image using the HRP form of this antibody.




Immunohistological and western blot analyses of HIF-1 α . (A) PAB liver stained by HIF-1 α (original magnification, $\times 20$). (B) Sham-operated control liver stained by HIF-1 α (original magnification, $\times 20$). (C) Enlarged view of the boxed area in the Image (A) (original magnification, $\times 100$). (D) Western blot analysis of HIF-1 α (mean \pm SEM), * $P < 0.05$. HIF-1 α : hypoxia-inducible factor-1 α ; PAB: pulmonary artery banding.




Immunocytochemistry/Immunofluorescence: Mouse Monoclonal HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Images of HIF-1 alpha localization (red) in CTR and CoCl₂-treated oAECs after 24 h of culture. Image from a verified customer review.



Western Blot: Mouse Monoclonal HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Image of HIF-1 alpha activation in AECs control cells (CTR) and after 24h of 10 and 50 μ M of CoCl₂-treatment; HIF-1 alpha time course at shorter CoCl₂ incubation times. Image from a verified customer review.




Western Blot: Mouse Monoclonal HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Image of HIF-1 alpha activation in AECs control cells (CTR) and after 24h of 10 and 50 μ M of CoCl₂-treatment; HIF-1 α time course at shorter CoCl₂ incubation times. Image from a verified customer review.




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunohistochemical detection of HIF-1 α protein in human tissues. The nature of the tissue is indicated on top of each figure. Original magnifications are as follows: A, \times 400; B, \times 400; C, \times 100; D, \times 100; E, \times 100; F, \times 100; G, \times 40; H, \times 40. Image collected & cropped by CiteAb from the following publication (<http://bmccgenet.biomedcentral.com/articles/10.1186/1471-2156-5-27>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.

G) Brain, glioblastoma




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunohistochemical detection of HIF-1 α protein in human tissues. The nature of the tissue is indicated on top of each figure. Original magnifications are as follows: A, \times 400; B, \times 400; C, \times 100; D, \times 100; E, \times 100; F, \times 100; G, \times 40; H, \times 40. Image collected & cropped by CiteAb from the following publication (<http://bmccgenet.biomedcentral.com/articles/10.1186/1471-2156-5-27>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunohistochemical detection of HIF-1 α protein in human tissues. The nature of the tissue is indicated on top of each figure. Original magnifications are as follows: A, \times 400; B, \times 400; C, \times 100; D, \times 100; E, \times 100; F, \times 100; G, \times 40; H, \times 40. Image collected & cropped by CiteAb from the following publication (<http://bmccgenet.biomedcentral.com/articles/10.1186/1471-2156-5-27>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.

D) Kidney, carcinoma




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunohistochemical detection of HIF-1 α protein in human tissues. The nature of the tissue is indicated on top of each figure. Original magnifications are as follows: A, $\times 400$; B, $\times 400$; C, $\times 100$; D, $\times 100$; E, $\times 100$; F, $\times 100$; G, $\times 40$; H, $\times 40$. Image collected & cropped by CiteAb from the following publication (<http://bmccgenet.biomedcentral.com/articles/10.1186/1471-2156-5-27>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.

E) Colon, normal




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunohistochemical detection of HIF-1 α protein in human tissues. The nature of the tissue is indicated on top of each figure. Original magnifications are as follows: A, $\times 400$; B, $\times 400$; C, $\times 100$; D, $\times 100$; E, $\times 100$; F, $\times 100$; G, $\times 40$; H, $\times 40$. Image collected & cropped by CiteAb from the following publication (<http://bmccgenet.biomedcentral.com/articles/10.1186/1471-2156-5-27>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.

H) Skin, melanoma




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunohistochemical detection of HIF-1 α protein in human tissues. The nature of the tissue is indicated on top of each figure. Original magnifications are as follows: A, $\times 400$; B, $\times 400$; C, $\times 100$; D, $\times 100$; E, $\times 100$; F, $\times 100$; G, $\times 40$; H, $\times 40$. Image collected & cropped by CiteAb from the following publication (<http://bmccgenet.biomedcentral.com/articles/10.1186/1471-2156-5-27>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.

F) Colon, adenocarcinoma




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunohistochemical detection of HIF-1 α protein in human tissues. The nature of the tissue is indicated on top of each figure. Original magnifications are as follows: A, $\times 400$; B, $\times 400$; C, $\times 100$; D, $\times 100$; E, $\times 100$; F, $\times 100$; G, $\times 40$; H, $\times 40$. Image collected & cropped by CiteAb from the following publication (<http://bmccgenet.biomedcentral.com/articles/10.1186/1471-2156-5-27>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.


C) Kidney, normal



Western Blot: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunohistological & western blot analyses of HIF-1 α . (A) PAB liver stained by HIF-1 α (original magnification, $\times 20$). (B) Sham-operated control liver stained by HIF-1 α (original magnification, $\times 20$). (C) Enlarged view of the boxed area in the Image (A) (original magnification, $\times 100$). (D) Western blot analysis of HIF-1 α (mean \pm SEM), * $P < 0.05$. HIF-1 α : hypoxia-inducible factor-1 α ; PAB: pulmonary artery banding. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/26863419>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunocytochemistry/ Immunofluorescence: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Analysis of CBF treatment in HCT116 implanted mouse models & tumour tissue samples. (a) Tumour size versus CBF treatment. The lowest tumour growth rate was found in i.p. group during CBF treatment. (b) Analysis of HIF-1 α mRNA level in tumour tissues. There was a significant upregulated HIF-1 α level in i.p. group. Results are means with standard errors from 4 replicates. (c) Inhibition of HIF-1 α nuclear translocation in the i.t. sample. CBF strongly inhibited HIF-1 α nuclear accumulation in the tumour of xenografts implanted with HCT116 cells. Scale bars equal 10 μ m. White arrows: HIF-1 α in the cytosol. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/23818933>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunohistochemical detection of HIF-1 α protein in human tissues. The nature of the tissue is indicated on top of each figure. Original magnifications are as follows: A, $\times 400$; B, $\times 400$; C, $\times 100$; D, $\times 100$; E, $\times 100$; F, $\times 100$; G, $\times 40$; H, $\times 40$. Image collected & cropped by CiteAb from the following publication (<http://bmccgenet.biomedcentral.com/articles/10.1186/1471-2156-5-27>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.

A) Lung, normal




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - CycT alleviates tumor hypoxia. The effects of CycT on the levels of exogenous hypoxia-marker pimonidazole labeling (A), the levels of hypoxia-inducible CA9 enzyme (B), & the levels of hypoxia-inducible factor HIF1 α (C) in orthotopic tumor xenografts. Scale bar: Montage, 1 mm; 10X, 20 μ m. Data are plotted as mean \pm SEM. For statistical analysis, the levels in treated tumors were compared to the levels in control tumors with a Welch 2-sample t-test. **p-value < 0.005 . IHC images are representative of 3 independent experiments. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/30723259>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Immunohistochemical detection of HIF-1 α protein in human tissues. The nature of the tissue is indicated on top of each figure. Original magnifications are as follows: A, $\times 400$; B, $\times 400$; C, $\times 100$; D, $\times 100$; E, $\times 100$; F, $\times 100$; G, $\times 40$; H, $\times 40$. Image collected & cropped by CiteAb from the following publication (<http://bmccgenet.biomedcentral.com/articles/10.1186/1471-2156-5-27>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.


B) Lung, carcinoma




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Schematic demonstration of the three HIF1 α antibodies used in this study & their epitopes (Figs. 2a & b). Ab1 is a polyclonal antibody raised against N-domain of wild type HIF1 α & does not recognize HIF1 α 1.2 due to its different N-terminal part (Fig. 2a). Ab2 & Ab3 are monoclonal & polyclonal antibodies, respectively, with epitopes in the common parts of HIF1 α & HIF1 α 1.2 (Fig.2a & b). Immunohistochemical analysis performed on thin adjacent sections of NE-differentiated prostate cancer using the HIF1 α antibodies Ab1 (Fig. 2c), Ab 2 (Fig. 2d), Ab 3 (Fig. 2e) & HIF1 β antibody (Fig. 2f). Immunopositivity was detected for HIF1 α with Ab2 & Ab3 while HIF1 α Ab1 produced no detectable staining. HIF1 β was also positive in adjacent section (Fig. 2f). Double staining of chromogranin A & androgen receptor antigens on adjacent sections (Fig. 2g) showed immunopositivity for chromogranin A (Fast red). Androgen receptor antibody (DAB, brown) produced no staining. Immunostaining of benign prostate tissue with HIF1 α Ab3 showed immunopositivity in NE-like cells of benign prostate tissue (Fig. 2h). In addition, HIF1 β antibody recognized benign NE-like cells in benign prostate hyperplasia (Fig. 2i). Double staining with HIF1 α Ab3 & HIF1 β (Fig. 2j) shows co-localization of the two proteins in NE-like cells of benign prostate hyperplasia (HIF1 α Ab3 red stain, HIF1 β brown stain). Panels a, b, c, d, e, f & g: 40 \times objective. Panels h, i & j: 60 \times objective. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/20663134>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunocytochemistry/ Immunofluorescence: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - GAD65/67-positive neurons expressed HIF-1 α under hypoxic conditions. A) HIF-1 α expression co-localized with GAD65/67-ir in neurons exposed to hypoxia when compared to normoxia (upper panel) or GAD65/67-negative neurons in hypoxia (bottom panel, open arrow). B) Quantification shows the percentage of HIF-1 α -expressing GAD65/67-positive neurons after hypoxia in vitro (mean \pm SD; from n = 6 cultures). C) In vivo immunostaining illustrates HIF-1 α -positive (bottom panel, solid arrow) & HIF-1 α -negative (bottom panel, open arrow) in GAD65/67-ir neurons in the ipsilateral region, whereas the contralateral region shows no HIF-1 α staining in GAD65/67-ir neurons (see Figure 1 for region selection). Scale bars, 10 μ m (A); 20 μ m (B). Image collected & cropped by CiteAb from the following publication (<https://actaneurocomms.biomedcentral.com/articles/10.1186/2051-5960-2-51>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunocytochemistry/ Immunofluorescence: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - GAD65/67-positive neurons expressed HIF-1 α under hypoxic conditions. A) HIF-1 α expression co-localized with GAD65/67-ir in neurons exposed to hypoxia when compared to normoxia (upper panel) or GAD65/67-negative neurons in hypoxia (bottom panel, open arrow). B) Quantification shows the percentage of HIF-1 α -expressing GAD65/67-positive neurons after hypoxia in vitro (mean \pm SD; from n = 6 cultures). C) In vivo immunostaining illustrates HIF-1 α -positive (bottom panel, solid arrow) & HIF-1 α -negative (bottom panel, open arrow) in GAD65/67-ir neurons in the ipsilateral region, whereas the contralateral region shows no HIF-1 α staining in GAD65/67-ir neurons (see Figure 1 for region selection). Scale bars, 10 μ m (A); 20 μ m (B). Image collected & cropped by CiteAb from the following publication (<https://actaneurocomms.biomedcentral.com/articles/10.1186/2051-5960-2-51>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Results of in situ hybridization & immunohistochemistry on thin adjacent section to detect expression of HIF-1 α 1.2 mRNA & HIF1 α protein in malignant & benign prostate tissue. In situ hybridization (antisense probe, Fig. 3a) & immunostaining with HIF1 α Ab2 (Fig. 3c) on thin adjacent sections of NE-differentiated prostate adenocarcinoma showed co-localization of HIF1 α 1.2 transcript & HIF-1 α protein. Incubation with sense probe did not generate any detectable hybridization signals (Fig. 3b). Both In situ hybridization (Fig. 3d antisense) & HIF-1 α Ab2 immunostaining (Fig. 3f) were negative in non-NE-differentiated prostate adenocarcinoma. In situ hybridization with sense probe performed on non-NE-differentiated prostate cancer (Fig. 3e) was negative. In situ hybridizing on benign prostate tissue showed HIF1 α 1.2 transcript in NE-like cells of benign prostate tissue (Fig. 3g, \square). The sense probe on thin adjacent section generated no signals (Fig. 3h, \square). Furthermore, co-localization of HIF1 α 1.2 transcript (Fig. 3i, $\square, \square, \square$) & HIF1 α protein, detected with HIF1 α Ab3 (Fig. 3j, $\square, \square, \square$) was also shown in NE-like cells of benign prostate tissue. Panels a, b, c, d, e, f, g, h, i, j: 40 \times objective. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/20663134>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - HX-induced Sirt1 expression in white matter OPCs requires HIF1 α . (a) HIF1 α stabilization of Sirt1 transcript expression in OPCs as revealed by representative RT-PCR represents higher level of Sirt1 mRNA in VHL cKO mice. GDPDH mRNA serves as a control. Mean \pm s.e.m., n=3 brains for each group. (b,c) Representative western blot demonstrates a transient increase of HIF1 α expression in HX white matter at P18 with no significant effect at P11 (P=0.7955) & P45 (P=0.7333). Histograms show mean \pm s.e.m. (d,e) Graphs represent the percentages of Sirt1+Ki67+ & NG2+Ki67+ cells after HX white matter in WT & HIF1 α KO mice. Number in parentheses within bar indicates number of samples (n=4 brains per group & per genotype; ****P<0.0001, one-way analysis of variance, Bonferroni post hoc test, mean \pm s.e.m.). (f,g) Western blot demonstrates no increase (P=0.8231) in Sirt1 & HIF1 α expression in white matter of Hif1 α KO mice. Actin was used as loading control (mean \pm s.e.m.; n=3 brains for each experiment, group & genotype). Image collected & cropped by CiteAb from the following publication (<https://www.nature.com/articles/ncomms13866>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - Results of in situ hybridization & immunohistochemistry on thin adjacent section to detect expression of HIF-1 α 1.2 mRNA & HIF1 α protein in malignant & benign prostate tissue. In situ hybridization (antisense probe, Fig. 3a) & immunostaining with HIF1 α Ab2 (Fig. 3c) on thin adjacent sections of NE-differentiated prostate adenocarcinoma showed co-localization of HIF1 α 1.2 transcript & HIF-1 α protein. Incubation with sense probe did not generate any detectable hybridization signals (Fig. 3b). Both In situ hybridization (Fig. 3d antisense) & HIF-1 α Ab2 immunostaining (Fig. 3f) were negative in non-NE-differentiated prostate adenocarcinoma. In situ hybridization with sense probe performed on non-NE-differentiated prostate cancer (Fig. 3e) was negative. In situ hybridizing on benign prostate tissue showed HIF1 α 1.2 transcript in NE-like cells of benign prostate tissue (Fig. 3g, \square). The sense probe on thin adjacent section generated no signals (Fig. 3h, \square). Furthermore, co-localization of HIF1 α 1.2 transcript (Fig. 3i, $\square, \square, \square$) & HIF1 α protein, detected with HIF1 α Ab3 (Fig. 3j, $\square, \square, \square$) was also shown in NE-like cells of benign prostate tissue. Panels a, b, c, d, e, f, g, h, i, j: 40 \times objective. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/20663134>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunocytochemistry/ Immunofluorescence: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - HIF-1 α expression in primary cortical neurons exposed to hypoxia/ischemia. A & A') Neurons were double-stained for HIF-1 α & MAP2 in the presence of 5 & 25 mM glucose with & without hypoxia. HIF-1 α expression in the somata was observed in cells with interneuron-like morphology after hypoxia. B) CoCl₂ (0.3 mM) induced HIF-1 α expression in cells with interneuron-like morphology. C) Quantification represents the increase in HIF-1 α -ir staining (mean \pm SD; 5-10 neurons quantified from each experiment, n = 3 experiments). D) In vivo brain slice shows a similar pattern of positive HIF-1 α -ir in round soma (open arrow) & negative in neurons with pyramidal-like morphology (solid arrow) in the ipsilateral side. *p < 0.05, compared with normoxia (25 mM glucose), #p < 0.05, compared with hypoxia (25 mM glucose). Scale bars, 20 μ m (A, A', B); 10 μ m (D). Image collected & cropped by CiteAb from the following publication (<https://actaneurocomms.biomedcentral.com/articles/10.1186/2051-5960-2-51>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: HIF-1 alpha Antibody (H1alpha67) [NB100-123] - HX-induced Sirt1 expression in white matter OPCs requires HIF1 α . (a) HIF1 α stabilization of Sirt1 transcript expression in OPCs as revealed by representative RT-PCR represents higher level of Sirt1 mRNA in VHL cKO mice. GDPDH mRNA serves as a control. Mean \pm s.e.m., n=3 brains for each group. (b,c) Representative western blot demonstrates a transient increase of HIF1 α expression in HX white matter at P18 with no significant effect at P11 (P=0.7955) & P45 (P=0.7333). Histograms show mean \pm s.e.m. (d,e) Graphs represent the percentages of Sirt1+Ki67+ & NG2+Ki67+ cells after HX white matter in WT & HIF1 α KO mice. Number in parentheses within bar indicates number of samples (n=4 brains per group & per genotype; **P<0.0001, one-way analysis of variance, Bonferroni post hoc test, mean \pm s.e.m.). (f,g) Western blot demonstrates no increase (P=0.8231) in Sirt1 & HIF1 α expression in white matter of Hif1 α KO mice. Actin was used as loading control (mean \pm s.e.m.; n=3 brains for each experiment, group & genotype). Image collected & cropped by CiteAb from the following publication (<https://www.nature.com/articles/ncomms13866>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.**



Publications

- Mandl M, Depping R. ARNT is a potential direct HIF-1 target gene in human Hep3B hepatocellular carcinoma cells. *Cancer Cell Int.* 2017-08-31 [PMID: 28855849]
- Nanduri J, Wang N, Wang BI, Prabhakar Nr Lysine demethylase KDM6B regulates HIF-1 alpha mediated systemic and cellular responses to intermittent hypoxia *Physiological genomics* 2021-07-23 [PMID: 34297635]
- Jenna Kerry, Erin J Specker, Morgan Mizzoni, Andrea Brumwell, Leslie Fell, Jenna Goodbrand, Michael N Rosen, James Uniacke Autophagy-dependent alternative splicing of ribosomal protein S24 produces a more stable isoform that aids in hypoxic cell survival. *FEBS letters* 2024-03-12 [PMID: 38281767]
- Yeh JL, Kuo CH, Shih PW et al. Xanthine derivative KMUP-1 ameliorates retinopathy *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* 2023-07-03 [PMID: 37406513]
- Kerry J, Specker E, Mizzoni M et al. Autophagy-dependent alternative splicing event produces a more stable ribosomal protein S24 isoform that aids in hypoxic cell survival *bioRxiv* 2023-09-26 (WB)
- Bharti A, Urs AB, Kumar P. Significance of HIF-1? Expression and LOXL-2 Localization in Progression of Oral Squamous Cell Carcinoma *Asian Pacific Journal of Cancer Prevention* 2021-02-01 [PMID: 33639646] (Immunohistochemistry, Immunohistochemistry-Paraffin)
- Li X, Ma TK, Wang M et al. YY1-induced upregulation of LncRNA-ARAP1-AS2 and ARAP1 promotes diabetic kidney fibrosis via aberrant glycolysis associated with EGFR/PKM2/HIF-1? pathway *Frontiers in Pharmacology* 2023-02-15 [PMID: 36874012]
- Segura S, Stolnicu S, Boros M et al. mTOR Pathway Activation Assessed by Immunohistochemistry in Cervical Biopsies of HPV-associated Endocervical Adenocarcinomas (HPVA): Correlation With Silva Invasion Patterns *Applied Immunohistochemistry & Molecular Morphology* 2021-08-01 [PMID: 33587450] (Immunohistochemistry)
- Kang HJ, Min BK, Choi WI et al. Pyruvate dehydrogenase kinase 1 and 2 deficiency reduces high-fat diet-induced hypertrophic obesity and inhibits the differentiation of preadipocytes into mature adipocytes *Experimental & Molecular Medicine* 2021-09-22 [PMID: 34552205] (Western Blot)
- Guo Y, Zhou J, Li X et al. The Association of Suppressed Hypoxia-Inducible Factor-1 Transactivation of Angiogenesis With Defective Recovery From Cerebral Ischemic Injury in Aged Rats *Frontiers in Aging Neuroscience* 2021-02-26 [PMID: 33716719]
- Heyer, V;Reina-San-Martin, B; Optimal AID expression and efficient immunoglobulin class switch recombination are dependent on the Hypoxia-Inducible Factor *European journal of immunology* 2023-05-04 [PMID: 37143384] (WB, Mouse)
- Pang X, Wang Z, Yin D, Zhang Z. Overexpression of hypoxia-inducible factor prolyl- hydroxylase attenuated by HCG-induced vascular endothelial growth factor expression in luteal cells. *Mol Med Rep* 2015-05-16 [PMID: 25975603]
- More publications at <http://www.novusbio.com/NB100-123>



Procedures

Western Blot protocol for HIF-1 alpha Antibody (NB100-123)

Western Blot Protocol

1. Resolve aliquots (25-30 ug) of induced nuclear protein extracts on a 4-20% Tris-HCl gel.
2. Transfer proteins to nitrocellulose membrane in 20 mM Tris-HCL (pH 8.0)/150 mM glycine/20% (vol/vol) methanol.
3. Block membrane for 1 hour with 1X western wash buffer containing 5% non-fat dry milk (NFDM).
4. Incubate membrane overnight at 4C in NB 100-123 diluted in 1X western wash/5% NFDM.
5. Wash with 1X western wash for 35 minutes at RT (1 X 15 minutes, 2 X 10 minutes).
6. Incubate membrane with HRP conjugated anti-mouse IgG for 1 hour (RT) in 1X western wash/5% NFDM.
7. Wash with 1X western wash for 35 minutes at RT (1 X 15 minutes, 2 X 10 minutes).
8. Drain membrane and place on saran wrap.
9. Using Amersham ECL Kit, mix equal volumes of two reagents. Pour over membrane (protein side facing up). Let solution sit on membrane for 15-20 seconds.
10. Drain membrane and place on new saran wrap.
11. Wrap up membrane and expose to film.
12. Develop accordingly.

10X Western wash: 24.2 g Tris, 80g NaCl, Tween-20 to 1%, pH 7.6 and QS to 4L.

Stripping buffer: 100 mM BME 2% SDS 62.5 mM Tris (pH 6.7)

To strip membrane: Incubate membrane in stripping buffer for 30 minutes at 56C. Wash membrane for 15 minutes with several change of 1X western wash.

Notes: If hypoxia treatment is not hypoxic enough (less than 2% oxygen to get an induction), signal will be absent. Also, if the harvest time is too slow or there are not enough protease inhibitors, etc., the induced protein will be rapidly lost as HIF-1alpha has a very short half-life.

Nuclear Extract Preparation Reference: Wang and Semenza. Purification and Characterization of Hypoxia-Inducible Factor. Journal of Biological Chemistry. 270(3): 1230-1237, 1995.

**This antibody has demonstrated varying results in Western blot applications. Product NB100-105 is recommended for most Western blot experiments.

Immunohistochemistry protocol for HIF-1 alpha Antibody (NB100-123)

Please see:

Primary Reference: Zhong, H., et al. Overexpression of Hypoxia-inducible Factor 1alpha in Common Human Cancers and Their Metastases. Cancer Research. 59: 5830-5835, 1999.

Immunocytochemistry/Immunofluorescence protocol for HIF-1 alpha Antibody (NB100-123)

1. Fix cells in 3% Paraformaldehyde in PBS for 15 minutes at room temperature, gently rocking.
2. Rinse cells 3 times for 5 minutes in PBS.
3. Block and permeabilize cells in 2% non-fat dry milk (NFDM) dissolved in PBS with 0.1% TX-100 overnight at 4C (covered to prevent evaporation).
4. Rinse cells 3 times for 5 minutes in PBS.
5. Dilute NB100-123 1:500 in dilution buffer [2% BSA in PBS with 0.01% TX-100].
6. Place cover slip upside down on a 50 ul drop of diluted antibody on parafilm, in humidity box.
7. Incubate for 1 hour at 37C.
8. Flip slips right side up in wells and rinse 3 times for 5 minutes each, in PBS.
9. In an amber microfuge tube, dilute secondary antibody (Cy3 anti-ms IgG) 1:500 in dilution buffer [2% BSA in PBS with 0.01% TX-100].
10. Place 800 ul of diluted secondary antibody in each well and make sure the fluid film covers over the cells on the slip. Alternatively, secondary antibody can be applied in the same manner as the primary (slip upside-down on drop of secondary that has been placed on a sheet of parafilm that is inside of a humidity box).
11. Incubate for 1 hour at 37C, in the dark.
12. Rinse cells at room temperature 4 times for 15 minutes each, in PBS, gently rocking.
13. Mount on frosted slides with AquaPoly Mount (Polysciences).
14. Refrigerate flat and covered.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-123

NBP2-36452	HeLa Hypoxic / Normoxic Cell Lysate
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP2-27231	Mouse IgG2b Isotype Control (MPC-11)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-123

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

