

Product Datasheet

Pleckstrin Antibody NB100-1343

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-1343

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-1343



NB100-1343**Pleckstrin Antibody**


Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA
Target Molecular Weight	40 kDa

Product Description	
Description	Novus Biologicals Goat Pleckstrin Antibody (NB100-1343) is a polyclonal antibody validated for use in IHC, WB, ELISA, Flow and ICC/IF. Anti-Pleckstrin Antibody: Cited in 2 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	5341
Gene Symbol	PLEK
Species	Human
Immunogen	Peptide with sequence C-EWIKAIQMASRTGK corresponding to C-Terminus according to NP_002655.2.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Peptide ELISA
Recommended Dilutions	Western Blot 0.01 - 0.3 ug/mL, Flow Cytometry 10 ug/mL, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 10 ug/mL, Immunohistochemistry-Paraffin 3.75 ug/mL, Peptide ELISA Detection limit 1:16000
Application Notes	WB: Approx. 45 kDa band observed in human peripheral blood mononucleocyte lysates (calculated MW of 40.1 kDa band according to NP_002655.2).

Images


Western Blot: Pleckstrin Antibody [NB100-1343] - Staining of Human Parathyroid lysate (A) and negative control HepG2 (B) cell lysate (35ug protein in RIPA buffer). Antibody at 0.01 ug/mL. Detected by chemiluminescence.




Immunocytochemistry/Immunofluorescence: Pleckstrin Antibody [NB100-1343] - Analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL), showing cytoplasmic/membrane and nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL).




Immunohistochemistry-Paraffin: Pleckstrin Antibody [NB100-1343] - Staining of paraffin embedded Human Spleen. Antibody at 3.75 ug/mL. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Flow Cytometry: Pleckstrin Antibody [NB100-1343] - Analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (1 ug/mL). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



Immunohistochemistry-Paraffin: Pleckstrin Antibody [NB100-1343] - Staining of paraffin embedded Human Vessel (platelets). Antibody at 3.75 ug/mL. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Publications

Tyers M, Rachubinski RA, Stewart MI et al. Molecular cloning and expression of the major protein kinase C substrate of platelets. *Nature* 1988-06-02 [PMID: 2897630]

Hunter RW, Mackintosh C, Hers I et al. Protein kinase C-mediated phosphorylation and activation of PDE3A regulate cAMP levels in human platelets. *J Biol Chem*. 2009-05-01 [PMID: 19261611]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950

Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada

Phone: 905.827.6400

Toll Free: 855.668.8722

Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15

Fax: (44) (0) 1235 533420

info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB100-1343

NBL1-14499	Pleckstrin Overexpression Lysate
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1343

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



