Product Datasheet

MDM2/HDM2 [p Ser185] Antibody [Biotin] NB100-1683B

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-1683B

Updated 7/11/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB100-1683B



NB100-1683B

MDM2/HDM2 [p Ser185] Antibody [Biotin]

Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Polyclonal Preservative 0.05% Sodium Azide Isotype IgG Conjugate Biotin Purity Immunogen affinity purified Buffer PBS Product Description Host Rabbit Gene ID 4193 Gene Symbol MDM2 Species Human, Mouse Reactivity Notes ABLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the phosphorylated form of the immunizing peptide. Reactivity occurs against the non-phosphorylated form of the immunizing peptide. Reactivity wouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody wis specific for the phosphorylated form of the immunizing peptide. Reactivity occurs against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against the mon-phosphorylated form of the manual peptide. Reactivity occurs against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against the non-phosphorylated form of the protein. Reactivity with non-phosphorylated form of the protein. Reactivity with non-phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 protein and the antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse MDM2/HDM2/HDM2/HDM2. (Uniprot: P23804)	MDM2/11DM2 [p GC1100] / Intibody [Diotin]	
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Polyclonal Preservative 0.05% Sodium Azide Isotype IgG Conjugate Biotin Purity Immunogen affinity purified Buffer PBS Product Description Host Rabbit Gene ID 4193 Gene Symbol MDM2 Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Product Information	
Storage Store at 4C in the dark. Clonality Polyclonal Preservative 0.05% Sodium Azide Isotype IgG Conjugate Biotin Purity Immunogen affinity purified Buffer PBS Product Description Host Rabbit Gene ID 4193 Gene Symbol MDM2 Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with hon-phosphorylated mouse MDM2/HDM2 is minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Unit Size	0.1 ml
Clonality Polyclonal Preservative 0.05% Sodium Azide Isotype IgG Conjugate Biotin Purity Immunogen affinity purified Buffer PBS Product Description Host Rabbit Gene ID 4193 Gene Symbol MDM2 Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 pS185 and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Concentration	·
Preservative Isotype IgG Conjugate Biotin Purity Immunogen affinity purified Buffer PBS Product Description Host Rabbit Gene ID 4193 Gene Symbol MDM2 Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Storage	Store at 4C in the dark.
Isotype	Clonality	Polyclonal
Conjugate Biotin Purity Immunogen affinity purified Buffer PBS Product Description Host Rabbit Gene ID 4193 Gene Symbol MDM2 Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Preservative	0.05% Sodium Azide
Purity Immunogen affinity purified Buffer PBS Product Description Host Rabbit Gene ID 4193 Gene Symbol MDM2 Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Isotype	IgG
Product Description Host Rabbit Gene ID 4193 Gene Symbol MDM2 Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Conjugate	Biotin
Product Description Host Rabbit Gene ID 4193 Gene Symbol MDM2 Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Purity	Immunogen affinity purified
Host Rabbit Gene ID 4193 Gene Symbol MDM2 Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Buffer	PBS
Gene Symbol MDM2 Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Product Description	
Gene Symbol MDM2 Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Host	Rabbit
Species Human, Mouse Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Gene ID	4193
Reactivity Notes A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Gene Symbol	MDM2
MDM2/HDM2 homologues from other sources. Specificity/Sensitivity This affinity-purified antibody is directed against the phosphorylated form of mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Species	Human, Mouse
mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity with MDM2/HDM2 homologues from other sources. Immunogen This affinity purified MDM2/HDM2 [p Ser185] Antibody was prepared from whole rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Reactivity Notes	
rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse	Specificity/Sensitivity	mouse MDM2/HDM2 protein at the pS185 residue. The resultant affinity purified antibody was then cross-adsorbed against the non-phosphorylated form of the immunizing peptide. Reactivity occurs against Mouse MDM2/HDM2 pS185 protein and the antibody is specific for the phosphorylated form of the protein. Reactivity with non-phosphorylated mouse MDM2/HDM2 is minimal by ELISA and western blot. A BLAST analysis was used to suggest minimal cross reactivity
	Immunogen	rabbit serum produced by repeated immunizations with a synthetic peptide corresponding to an internal region near aa 175-200 of mouse



Notes

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

Product Application Details	
Applications	Western Blot, ELISA, Immunoprecipitation
Recommended Dilutions	Western Blot, ELISA, Immunoprecipitation
Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966 novus@novusbio.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: technical@novusbio.com

Orders: orders@novusbio.com General: novus@novusbio.com

Products Related to NB100-1683B

NBP2-24891B Rabbit IgG Isotype Control [Biotin]

DYC1244-2 MDM2/HDM2 [Biotin]

NB600-302 c-Myc Antibody (9E10) - BSA Free

NB200-106 p19ARF/CDKN2A Antibody - BSA Free

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1683B

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

