

Product Datasheet

Glutathione Peroxidase 2/GPX2 Antibody NB100-1888

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 6

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-1888

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-1888



NB100-1888

Glutathione Peroxidase 2/GPX2 Antibody


Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA

Product Description	
Description	Novus Biologicals Goat Glutathione Peroxidase 2/GPX2 Antibody (NB100-1888) is a polyclonal antibody validated for use in IHC, WB and ELISA. Anti-Glutathione Peroxidase 2/GPX2 Antibody: Cited in 6 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	2877
Gene Symbol	GPX2
Species	Human, Mouse, Rat
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 19151108); Rat reactivity reported in scientific literature (PMID: 18056462).
Immunogen	Peptide with sequence C-SLDGEKVDFN corresponding to internal region according to NP_002074.2.

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Peptide ELISA
Recommended Dilutions	Western Blot 0.5 - 2 ug/ml, Immunohistochemistry 1:10 - 1:500, Immunohistochemistry-Paraffin, Peptide ELISA Detection limit 1:32000
Application Notes	WB: Approx. 24 kDa band observed in human liver lysates (calculated MW of 21.9 kDa band according to NP_002074.2). Use in IHC-P reactivity reported in (PMID: 24562575).

Images

Western Blot: Glutathione Peroxidase 2/GPX2 Antibody [NB100-1888] - Staining (0.5ug/ml) of Human Liver lysate (RIPA buffer, 30ug total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.



Publications

Doi K, Fujioka M, Sokuza Y et al. Chemopreventive Action by Ethanol-extracted Brazilian Green Propolis on Post-initiation Phase of Inflammation-associated Rat Colon Tumorigenesis. *In Vivo*. 2017-03-30 [PMID: 28358699] (IF/IHC, Mouse)

Esworthy RS, Yang L, Frankel PH, Chu FF. Epithelium-specific glutathione peroxidase, Gpx2, is involved in the prevention of intestinal inflammation in selenium-deficient mice. *J Nutr* 2005-04-01 [PMID: 15795427]

Naiki T, Naiki-Ito A, Asamoto M et al. GPX2 overexpression is involved in cell proliferation and prognosis of castration resistant prostate cancer. *Carcinogenesis*. 2014-02-22 [PMID: 24562575] (IHC-P, WB, Rat, Human)

Details:

Rat castration-resistant prostate cancer model: Fig 1A [IHC(P), Fig 1B (WB); LnCAP and PC3 human prostate carcinoma cells: Fig 1B (WB); Human prostatectomy normal and cancerous specimens: Figs 4 & 5 [IHC (P)]. sections of the tissues were incubated with 1

Naiki-Ito A, Asamoto M, Hokaiwado N et al. Gpx2 is an overexpressed gene in rat breast cancers induced by three different chemical carcinogens. *Cancer Res*. 2007-12-01 [PMID: 18056462] (WB, IHC-P, Rat)

Details:

WB: rat normal mammary and carcinogen-induced mammary cancer (Fig 2B) IHC (paraffin): rat normal mammary tissue and carcinogen-induced mammary cancer (Fig 2C) and human normal breast and breast cancer (Fig 3).

McGrath-Morrow S, Lauer T, Yee M et al. Nrf2 increases survival and attenuates alveolar growth inhibition in neonatal mice exposed to hyperoxia. *Am J Physiol Lung Cell Mol Physiol*. 2009-04-01 [PMID: 19151108] (WB, Mouse)

Details:

Newborn mouse lung tissue: WB (Fig 4B, C); IHC (paraffin): Fig 4D.

Roland-Zejly L, Moisan V, St-Pierre I, Bilodeau JF. Altered placental glutathione peroxidase mRNA expression in preeclampsia according to the presence or absence of labor Placenta 2011-02-01 [PMID: 21145108] (IF/IHC, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-1888

NB820-59232	Human Liver Whole Tissue Lysate (Adult Whole Normal)
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1888

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



