

Product Datasheet

Protein Disulfide Isomerase/P4HB Antibody NB100-1921

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 2 Publications: 6

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-1921

Updated 10/23/2024 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-1921




NB100-1921**Protein Disulfide Isomerase/P4HB Antibody**


Product Information	
Unit Size	0.1 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	No Preservative
Isotype	IgG
Purity	Unpurified
Buffer	Whole antisera
Product Description	
Host	Rabbit
Gene ID	5034
Gene Symbol	P4HB
Species	Human, Mouse, Rat, Bovine, Primate
Marker	Endoplasmic Reticulum Marker
Immunogen	Purified full length P4hb isolated from rat liver.
Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:10000-1:20000, Immunohistochemistry 1:100-1:1000, Immunocytochemistry/ Immunofluorescence 1:5000-1:20000, Immunoprecipitation 1-3 ug, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen 1:100-1:1000
Application Notes	This P4hb antibody is useful for Immunocytochemistry/Immunofluorescence, Immunohistochemistry frozen and paraffin embedded sections, Immunoprecipitation and Western Blot.

Images


Normal human brain.



ICC/IF Confocal analysis of 3T3 cells using P4HB antibody (NB100-1921, 1:5). An Alexa Fluor 488-conjugated Goat to rabbit IgG was used as secondary antibody (green). Actin filaments were labeled with Alexa Fluor 568 phalloidin (red). DAPI was used to stain the cell nuclei (blue).



IHC analysis of a formalin fixed paraffin-embedded (FFPE) human brain using 1:250 conc. of P4HB antibody on a Bond Rx autostainer (Leica Biosystems). The assay involved 20 minutes of heat induced antigen retrieval (HIER) using 10mM sodium citrate buffer (pH 6.0) and endogenous peroxidase quenching with peroxide block. The sections were incubated with primary antibody for 30 minutes and Bond Polymer Refine Detection (Leica Biosystems) with DAB was used for signal development followed by counterstaining with hematoxylin. Whole slide scanning and capturing of representative images was performed using Aperio AT2 (Leica Biosystems). Cytoplasmic staining of P4HB in neurons was observed. Staining was performed by Histowiz.



P4HB staining in Hela cells detected with a Dylight 488 labeled secondary antibody (Green) with Hoechst 33258 nuclear counterstain (Blue).



Publications

Oleinik N, Kim J, Roth BM et al. Mitochondrial protein import is regulated by p17/PERMIT to mediate lipid metabolism and cellular stress Sci Adv 2019-09-01 [PMID: 31535025] (WB, Human)

Gamberucci A, Marcolongo P, Nemeth CE et al. GLUT10-Lacking in Arterial Tortuosity Syndrome-Is Localized to the Endoplasmic Reticulum of Human Fibroblasts Int J Mol Sci 2017-08-22 [PMID: 28829359] (ICC/IF, Human)

Omachi K, Kamura M, Teramoto K et al. A Split-Luciferase-Based Trimer Formation Assay as a High-throughput Screening Platform for Therapeutics in Alport Syndrome. Cell Chem Biol. 2018-02-21 [PMID: 29526710] (WB)

Jun HS, Lee YM, Song K et al. G-CSF improves murine G6PC3-deficient neutrophil function by modulating apoptosis and energy homeostasis Blood 2011-04-07 [PMID: 21292774]

Azad AK, Torrelles JB, Schlesinger LS. Mutation in the DC-SIGN cytoplasmic triacidic cluster motif markedly attenuates receptor activity for phagocytosis and endocytosis of mannose-containing ligands by human myeloid cells. J Leukoc Biol;84(6):1594-603. 2008-12-01 [PMID: 18772280] (ICC/IF, Human)

Safran, M et al. Thyroid hormone-dependent redistribution of the 55-kilodalton monomer of protein disulfide isomerase in cultured glial cells. Endocrinology;131(5):2413-8. 1992-11-01 [PMID: 1425439]





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-1921

NB820-60576	Human Brain Frontal Lobe Membrane Tissue Lysate (Adult Membrane Normal)
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1921

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

