

Product Datasheet

HSP90 Antibody (AC88) - BSA Free NB100-1972

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 7

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-1972

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-1972




NB100-1972

HSP90 Antibody (AC88) - BSA Free


Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	AC88
Preservative	0.09% Sodium Azide
Isotype	IgG1
Purity	Protein G purified
Buffer	PBS and 50% Glycerol
Target Molecular Weight	90 kDa
Product Description	
Host	Mouse
Gene ID	3320
Gene Symbol	HSP90AA1
Species	Human, Mouse, Rat, Porcine, Bacteria, Canine, Chicken, Fish, Guinea Pig, Primate, Rabbit, Sheep
Reactivity Notes	Recognizes Beluga, <i>C. elegans</i> , <i>Fusobacterium nucleatum</i> , <i>Mycobacterium leprae</i> , Mussel, Scallop and Water Mold HSP90. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions.
Specificity/Sensitivity	Recognizes both Hsp90 alpha and beta. It reacts with the rodent glucocorticoid receptor and with proliferation potential proteins (P2PS) with apparent molecular masses of 30-40 kDa.
Immunogen	<i>Achlya ambisexualis</i> (water mold) Hsp90.
Product Application Details	
Applications	Western Blot, Simple Western, Flow Cytometry, Flow (Intracellular), Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1 ug/ml, Simple Western, Flow Cytometry, Immunohistochemistry 1:10-1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1:10-1:500, Flow (Intracellular)
Application Notes	Use in ICC/IF reported in scientific literature (PMID: 30373228).

Images


Western Blot: HSP90 Antibody (AC88) [NB100-1972] - Analysis of HSP90, mAb (AC88): Lane 1: MW Marker; Lane 2: HSP90 (human), (native) ; Lane 3: HSP90 beta, (human) (recombinant); Lane 4: HSP90a (human), (recombinant); Lane 5: HeLa (heat shocked); Lane 6: 3T3 (heat shocked); Lane 7: PC-12 (heat shocked) ; Lane 8: CHO-K1 (heat shocked).




Immunohistochemistry: HSP90 Antibody (AC88) [NB100-1972] - Analysis of human spleen tissue stained with HSP90, mAb (AC88) at 10 ug/ml.




Flow (Intracellular): HSP90 Antibody (AC88) [NB100-1972] - An intracellular stain was performed on HeLa cells with HSP90 Antibody (AC88) NB100-1972PE (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to phycoerythrin. Image using the PE format of this antibody.




Flow Cytometry: HSP90 Antibody (AC88) [NB100-1972] - Analysis using the DyLight 488 conjugate of NB100-1972. Staining of 106 Jurkat cells using HSP90 mAb (AC88), DyLight 488 Conjugate at a concentration of 50ug/ml.




Flow Cytometry: HSP90 Antibody (AC88) [NB100-1972] - Analysis using the Alexa Fluor 405 conjugated HSP90 antibody, NB100-1972AF405 (purple) and fluorescence minus one control (orange). Staining of HSP90 in Primary CD4+ T-cells derived from human peripheral blood using 1uL of antibody per 1X10e5 cells. Image from verified customer review.



Flow Cytometry: HSP90 Antibody (AC88) [NB100-1972] - Analysis using the PE conjugate of NB100-1972. Staining of 10^6 Jurkat cells using HSP90 mAb (AC88), R-Phycoerythrin Conjugate at a concentration of 10ug/mL




Flow Cytometry: HSP90 Antibody (AC88) [NB100-1972] - An intracellular stain was performed on HeLa cells with HSP90 Antibody (AC88) NB100-1972AF647 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 2.5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 647.



Copyright © 2018 Novus Biologicals

Flow Cytometry: HSP90 Antibody (AC88) [NB100-1972] - An intracellular stain was performed on Jurkat cells with HSP90 Antibody (AC88) NB100-1972AF647 (blue) and a matched isotype control (orange). Cells were fixed with 4% PFA and then permeabilized with 0.1% saponin. Cells were incubated in an antibody dilution of 5 ug/mL for 30 minutes at room temperature. Both antibodies were conjugated to Alexa Fluor 647.



Copyright © 2018 Novus Biologicals

Publications

Onay Ucar E, Sengelen A, Mertoglu Kamali E Hsp27, Hsp60, Hsp70, or Hsp90 depletion enhances the antitumor effects of resveratrol via oxidative and ER stress response in human glioblastoma cells *Biochemical Pharmacology* 2023-02-01 [PMID: 36603687] (WB, Human)

Details:

1:1000 WB dilution

Panagioti E, Kurokawa C, Viker K Et Al. Immunostimulatory bacterial antigen-armed oncolytic measles virotherapy significantly increases the potency of anti-PD1 checkpoint therapy *The Journal of clinical investigation* 2021-07-01 [PMID: 34196308] (FLOW)

Freund E, Miebach L, Stope MB, Bekeschus S Hypochlorous acid selectively promotes toxicity and the expression of danger signals in human abdominal cancer cells *Oncology reports* 2021-05-01 [PMID: 33760187] (FLOW, Human)

Mohamed H, Clemen R, Freund E et al. Non-thermal plasma modulates cellular markers associated with immunogenicity in a model of latent HIV-1 infection *PloS one* 2021-03-01 [PMID: 33647028]

Bekeschus S, Clemen R, Niebner F et al. Medical Gas Plasma Jet Technology Targets Murine Melanoma in an Immunogenic Fashion *Adv. Sci.* 2020-03-30 [PMID: 32440479]

Bekeschus S, Freund E, Wende k et al. Hmox1 Upregulation Is a Mutual Marker in Human Tumor Cells Exposed to Physical Plasma-Derived Oxidants. *Antioxidants (Basel)*. 2018-10-27 [PMID: 30373228] (ICC/IF, Human)

Kelly RL, Geoghegan JC, Feldman J et al. Chaperone proteins as single component reagents to assess antibody nonspecificity *MAbs* 2017-07-26 [PMID: 28745541] (Mouse)





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-1972

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)
NBC1-18362	Recombinant Human HSP90 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-1972

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

