

# Product Datasheet

## Blooms Syndrome Protein Blm Antibody - BSA Free NB100-214

Unit Size: 0.1 ml

Store at 4C. Do not freeze.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Publications: 8**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB100-214](http://www.novusbio.com/NB100-214)

Updated 2/21/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB100-214](http://www.novusbio.com/reviews/destination/NB100-214)



**NB100-214**

Blooms Syndrome Protein Blm Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)


Product Description	
Host	Rabbit
Gene ID	641
Gene Symbol	BLM
Species	Human, Mouse
Immunogen	The immunogen recognized by this antibody maps to a region between residue 1425 and the C-terminus (residue 1477) of human Bloom Syndrome using the numbering given in entry NP_000048.1 (GenelD 641).

Product Application Details	
Applications	Western Blot, Immunoprecipitation
Recommended Dilutions	Western Blot 1:2000-1:10000, Immunoprecipitation 2 to 10 ug/mg of lysate
Application Notes	Optimal dilutions should be determined by the end user.




## Images


**Western Blot: Blooms Syndrome Protein BLM Antibody [NB100-214] -**  
**Samples:** Whole cell lysate (15 ug) from HEK293T, K-562, Jurkat, HeLa, and A-549 cells prepared using NETN lysis buffer. **Antibody:** Affinity purified rabbit anti-BLM antibody NB100-214 (lot 4) used for WB at 0.1 ug/ml. **Detection:** Chemiluminescence with an exposure time of 30 seconds.



**Western Blot: Blooms Syndrome Protein BLM Antibody [NB100-214] -**  
**Samples:** Whole cell lysate (50 ug) from TCMK-1 and BIH 3T3 cells prepared using NETN lysis buffer. **Antibody:** Affinity purified rabbit anti-BLM antibody NB100-214 (lot 4) used for WB at 0.1 ug/ml. **Detection:** Chemiluminescence with an exposure time of 75 seconds.



**Immunoprecipitation: Blooms Syndrome Protein BLM Antibody [NB100-214] -**  
**Samples:** Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HEK293T cells prepared using NETN lysis buffer. **Antibodies:** Affinity purified rabbit anti-BLM antibody (NB100-214, lot 4) used for IP at 6 ug per reaction. BLM was also immunoprecipitated by a previous lot of this antibody (lot 3). For blotting immunoprecipitated BLM, NB100-214 was used at 0.1 ug/ml. **Detection:** Chemiluminescence with an exposure time of 10 seconds.



## Publications

Root H, Larsen A, Komosa M et al. FANCD2 limits BLM-dependent telomere instability in the alternative lengthening of telomeres pathway Hum. Mol. Genet. 2016-07-17 [PMID: 27427384]

Petrillo C, Barroca V, Ribeiro J et al. shani mutation in mouse affects splicing of Spata22 and leads to impaired meiotic recombination Chromosoma 2020-05-11 [PMID: 32388826]

Yoshimura Y Increased error-free DNA repair gene expression through reprogramming in human iPS cells Regenerative Therapy 2019-06-28 [PMID: 31304203] (WB, Human)

Nicholls TJ, Nadalutti CA, Motori E et al. Topoisomerase 3a Is Required for Decatenation and Segregation of Human mtDNA. Mol. Cell 2017-12-14 [PMID: 29290614] (Human)

Srivastava V, Modi P, Tripathi V et al. BLM helicase stimulates the ATPase and chromatin-remodeling activities of RAD54. J Cell Sci 2009-09-01 [PMID: 19671661] (Human)

Murfuni I, Nicolai S, Baldari S et al. The WRN and MUS81 proteins limit cell death and genome instability following oncogene activation Oncogene 2012-03-12 [PMID: 22410776]

Nair RR, Emmons MF, Cress AE et al. HYD1-induced increase in reactive oxygen species leads to autophagy necrotic cell death in multiple myeloma cells. Mol Cancer Ther;8(8):2441-2451. 2009-01-01 [PMID: 19671765]

Tripathi et al. BLM helicase dependent independent roles of 53BP1 during replication st. J Cell Biol 178:9. 2007-01-01 [PMID: 17591918]





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

[www.novusbio.com](http://www.novusbio.com)  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NB100-214**

---

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
H00000641-Q01-10ug	Recombinant Human Blooms Syndrome Protein Blm GST (N-Term) Protein

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB100-214](http://www.novusbio.com/reviews/submit/NB100-214)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

