## **Product Datasheet**

# RPA70 Antibody - BSA Free NB100-2204

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

**Publications: 7** 

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-2204

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB100-2204



## NB100-2204

RPA70 Antibody - BSA Free


Product Information		
Unit Size	100 ul	
Concentration	1.0 mg/ml	
Storage	Store at 4C. Do not freeze.	
Clonality	Polyclonal	
Preservative	0.09% Sodium Azide	
Isotype	IgG	
Purity	Immunogen affinity purified	
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)	
Due dont Description		

Product Description	
Description	Novus Biologicals Rabbit RPA70 Antibody - BSA Free (NB100-2204) is a polyclonal antibody validated for use in IHC, WB and IP. Anti-RPA70 Antibody: Cited in 7 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	6117
Gene Symbol	RPA1
Species	Human, Mouse
Reactivity Notes	Orangutan (100%).
Immunogen	The immunogen recognized by this antibody maps to a region between residues 350 and 400 of human replication protein A1, 70kDa using the numbering given in entry NP_002936.1 (GeneID 6117).

<b>Product Application Details</b>	3
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:2000-1:10000, Immunohistochemistry 1:1000-1:5000, Immunoprecipitation 2-10 ug/mg lysate, Immunohistochemistry-Paraffin 1:1000-1:5000
Application Notes	Immunohistochemistry 1:1,000 - 1:5,000. Epitope retrieval with Tris-EDTA pH 9.0 is recommended for FFPE tissue sections.


## **Images**

Western Blot: RPA70 Antibody [NB100-2204] - Whole cell lysate (50 ug) from NIH3T3 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit antiRPA70 antibody used for WB at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.







Immunohistochemistry-Paraffin: RPA70 Antibody [NB100-2204] - Human ovarian carcinoma. Antibodies: Affinity purified rabbit anti- RPA70 Cat. No. used at a dilution of 1:1,000 (1 ug/ml) (left) and used at a dilution of 1:5,000 (0.2ug/ml (right). Detection: DAB




Western Blot: RPA70 Antibody [NB100-2204] - Detection of Human RPA70 by Western Blot. Samples: Whole cell lysate (5, 15 and 50 ug) from HeLa and (15 and 50 ug) from 293T cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-RPA70 antibody NB100-2204 used for WB at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.



Immunohistochemistry-Paraffin: RPA70 Antibody [NB100-2204] - Serial sections of mouse teratoma. Antibodies: Affinity purified rabbit anti-RPA70 Lot2 used at a dilution of 1:1,000 (1 ug/ml) (left) and Lot3 used at a dilution of 1:5,000 (0.2ug/ml) (right). Detection: DAB



Immunoprecipitation: RPA70 Antibody [NB100-2204] - Detection of human RPA70 by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-RPA70 antibody NB100-2204 (lot NB100-2204-3) used for IP at 6 ug per reaction. RPA70 was also immunoprecipitated by a previous lot of this antibody (lot NB100-2204-2). For blotting immunoprecipitated RPA70, NB100-2204 was used at 1 ug/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.



#### **Publications**

Chatzidoukaki O, Stratigi K, Goulielmaki E Et al. R-loops trigger the release of cytoplasmic ssDNAs leading to chronic inflammation upon DNA damage Sci Adv 2021-11-19 [PMID: 34797720] (WB, Mouse)

#### Details:

Citation using the DyLight 550 version of this antibody.

Lindsey-Boltz LA, Reardon JT, Wold MS et al. In vitro analysis of the role of replication protein A (RPA) and RPA phosphorylation in ATR-mediated checkpoint signaling. J Biol Chem 2012-10-01 [PMID: 22948311]

Kemp MG, Reardon JT, Lindsey-Boltz LA et al. Mechanism of release and fate of excised oligonucleotides during nucleotide excision repair. J Biol Chem 2012-06-01 [PMID: 22573372]

Fisher LA, Bessho M, Wakasugi M et al. Role of interaction of XPF with RPA in nucleotide excision repair. J Mol Biol 2011-10-01 [PMID: 21875596]

Kemp MG, Lindsey-Boltz LA, Sancar A et al. The DNA damage response kinases DNA-dependent protein kinase (DNA-PK) and ataxia telangiectasia mutated (ATM) Are stimulated by bulky adduct-containing DNA. J Biol Chem 2011-06-01 [PMID: 21487018]

Thangavel S, Mendoza-Maldonado R, Tissino E et al. Human RECQ1 and RECQ4 helicases play distinct roles in DNA replication initiation. Mol Cell Biol 2010-03-01 [PMID: 20065033]

Robison JG, Elliott J, Dixon K et al. Replication protein A and the Mre11.Rad50.Nbs1 complex co-localize and interact at sites of stalled replication forks. J Biol Chem 2004-08-01 [PMID: 15180989]





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

## **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

## **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NB100-2204**

NBL1-15488 RPA70 Overexpression Lysate

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-2204

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

