

# Product Datasheet

## Jumonji/JARID2 Antibody - BSA Free NB100-2214

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Reviews: 1 Publications: 26**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB100-2214](http://www.novusbio.com/NB100-2214)

Updated 4/13/2025 v.20.1

**Earn rewards for product  
reviews and publications.**

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB100-2214](http://www.novusbio.com/reviews/destination/NB100-2214)



**NB100-2214**


Jumonji/JARID2 Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Glycine and 0.15M NaCl
Product Description	
Host	Rabbit
Gene ID	3720
Gene Symbol	JARID2
Species	Human, Mouse, Rabbit
Reactivity Notes	Human reactivity reported in scientific literature (PMID: 24074864)
Immunogen	A synthetic peptide made to an N-terminal portion of the human JARID2 protein sequence (between residues 1-100). [UniProt# Q92833]
Product Application Details	
Applications	Western Blot, Simple Western, Chromatin Immunoprecipitation, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation, Chromatin Immunoprecipitation (ChIP), Knockout Validated
Recommended Dilutions	Western Blot 1:500, Simple Western 1:10, Chromatin Immunoprecipitation 1:10-1:500. Use reported in scientific literature (PMID 22396653), Immunohistochemistry 1:200, Immunocytochemistry/ Immunofluorescence 1:1000, Immunoprecipitation 1:10-1:500. Use reported in scientific literature, Immunohistochemistry-Paraffin 1:200, Chromatin Immunoprecipitation (ChIP) 1:10-1:500, Knockout Validated
Application Notes	<p>By Western blot a band is seen at ~140 kDa. Another band is seen &gt;200 kDa and a faint band may be seen ~80 kDa. In ICC/IF, nuclear staining was observed in HeLa cells. In IHC-P, staining was observed in the nucleus of mouse brain. Prior to immunostaining paraffin tissues, antigen retrieval with sodium citrate buffer (pH 6.0) is recommended.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point. See <a href="#">Simple Western Antibody Database</a> for Simple Western validation: Tested in Human Brain lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:10, apparent MW was 128 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.</p>




## Images


**Western Blot: Jumonji/JARID2 Antibody [NB100-2214] - Jumonji/JARID2 N-terminus mediates direct interaction with H2AK119u1-modified nucleosomes.** Immunoblot of nucleosomal pull downs from nuclear extract. Total nuclear extract (input) and pull down using streptavidin beads, unmodified nucleosomes or nucleosomes modified by H2AK119u1/H2AK119u1(I44A) were probed with the indicated antibodies using extracts from WT cells (left) or Jumonji/JARID2 KO mESCs (right). Image collected and cropped by CiteAb from the following publication (<https://www.nature.com/articles/ncomms13661>) licensed under a CC-BY license.




**Immunocytochemistry/Immunofluorescence: Jumonji/JARID2 Antibody [NB100-2214] - JARID2 antibody was tested in HeLa cells with Dylight 488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and Dylight 550 (red).**




**Western Blot: Jumonji/JARID2 Antibody [NB100-2214] - Jarid2 expression in naive and activated (overnight) murine B cells.** This image was submitted via customer Review.




**Immunohistochemistry-Paraffin: Jumonji/JARID2 Antibody [NB100-2214] - Tested in mouse brain using DAB with hematoxylin counterstain.**




Western Blot: Jumonji/JARID2 Antibody [NB100-2214] - Detection of JARID2 protein in E16 mouse brain lysate using 2ug/ml of NB100-2214.



Simple Western: Jumonji/JARID2 Antibody [NB100-2214] - Simple Western lane view shows a specific band for JARID2 in 0.5 mg/ml of Human Brain lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



Western Blot: Jumonji/JARID2 Antibody [NB100-2214] - JARID2 N-terminus mediates direct interaction with H2AK119u1-modified nucleosomes(a) Immunoblot of nucleosomal pull downs from nuclear extract. Total nuclear extract (input) & pull down using streptavidin beads, unmodified nucleosomes or nucleosomes modified by H2AK119u1/H2AK119u1(I44A) were probed with the indicated antibodies using extracts from WT cells (left) or JARID2 KO mESCs (right). Full blots are shown in Supplementary Fig. 9. (b) JARID2 1–530 binds to H2AK119u1-modified mononucleosomes (yellow) more tightly than unmodified (blue) or H2AK199u1(I44A) (red) as monitored by biolayer interferometry (n=3). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/27892467>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



## Publications

Arecco N, Mocavini I, Blanco E, Ballaré C et Al. Alternative splicing decouples local from global PRC2 activity Mol Cell 2024-03-07 [PMID: 38452766]

Takeshi Suzuki MEG3 long noncoding RNA contributes to the epigenetic regulation of epithelial-mesenchymal transition in lung cancer cell lines J. Biol. Chem., 2016-11-16;0(0):. 2016-11-16 [PMID: 27852821]

Matteo Perino, Guido van Mierlo, Chet Loh, Sandra M.T. Wardle, Dick W. Zijlmans, Hendrik Marks, Gert Jan C. Veenstra Two Functional Axes of Feedback-Enforced PRC2 Recruitment in Mouse Embryonic Stem Cells Stem Cell Reports 2020-08-06 [PMID: 32763159]

Kate Skehan, Matthew Richardson, Laura M O'Connor, Samuel Dickson, Kate Martin, Geetha Govindarajulu, Swetha Sridharan Viscous Aqueous Gel Illustrating Natural Anatomy: The VAGINA method in gynaecological MRI simulation. Journal of medical radiation sciences 2023-08-24 [PMID: 37621131]

Nicola Reynolds, Mali Salmon-Divon, Heidi Dvinge, Antony Hynes-Allen, Gayan Balasooriya, Donna Leaford, Axel Behrens, Paul Bertone, Brian Hendrich NuRD-mediated deacetylation of H3K27 facilitates recruitment of Polycomb Repressive Complex 2 to direct gene repression The EMBO Journal 2012-02-01 [PMID: 22139358]

Liu W, Zeng Y, Hao X et al. JARID2 coordinates with the NuRD complex to facilitate breast tumorigenesis through response to adipocyte-derived leptin Cancer communications (London, England) 2023-09-01 [PMID: 37658635] (ICC/IF, IHC-P, WB, ChIP, Human)

Kadomatsu T, Hara C, Kurahashi R et al. ANGPTL2-mediated epigenetic repression of MHC-I in tumor cells accelerates tumor immune evasion Molecular oncology 2023-07-15 [PMID: 37452654] (IHC-Fr, Mouse)

Hickey GJ, Wike CL, Nie X et al. Establishment of developmental gene silencing by ordered polycomb complex recruitment in early zebrafish embryos eLife 2022-01-04 [PMID: 34982026]

Sun Z, Tang Y, Zhang Y Et al. Joint single-cell multiomic analysis in Wnt3a induced asymmetric stem cell division Nature communications 2021-10-12 [PMID: 34642323] (WB, Mouse)

Jain P, Ballare C, Blanco E et Al. PHF19 mediated regulation of proliferation and invasiveness in prostate cancer cells Elife 2020-03-10 [PMID: 32155117] (WB, Human)

Yen YP, Hsieh WF, Tsai YY et al. Dlk1-Dio3 locus-derived lncRNAs perpetuate postmitotic motor neuron cell fate and subtype identity Elife. 2018-10-11 [PMID: 30311912] (WB, IP, Mouse)

Kumar R, Evans T Activation-Induced Cytidine Deaminase Regulates Fibroblast Growth Factor/Extracellular Signal-Regulated Kinases Signaling To Achieve the Naive Pluripotent State During Reprogramming Stem Cells 2019-04-25 [PMID: 31021461] (WB, Mouse)

More publications at <http://www.novusbio.com/NB100-2214>





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NB100-2214**

---

NB100-2214PEP	Jumonji/JARID2 Antibody Blocking Peptide
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB100-2214](http://www.novusbio.com/reviews/submit/NB100-2214)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

