

Product Datasheet

u-Plasminogen Activator/Urokinase Antibody NB100-2419

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-2419

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-2419



NB100-2419

u-Plasminogen Activator/Urokinase Antibody

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris (pH 7.3), 0.5% BSA


Product Description	
Host	Goat
Gene ID	5328
Gene Symbol	PLAU
Species	Human
Specificity/Sensitivity	This antibody is expected to recognize both reported isoforms.
Immunogen	Peptide with sequence C-HFLPWIRSHTKEEN corresponding to C-Terminus according to NP_002649.1, NP_001138503.1.

Product Application Details	
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin, Peptide ELISA
Recommended Dilutions	Immunohistochemistry 8 ug/mL, Immunohistochemistry-Paraffin 8 ug/mL, Peptide ELISA Detection limit 1:6000



Images

Immunohistochemistry-Paraffin: u-Plasminogen Activator/Urokinase Antibody [NB100-2419] - Negative Control showing staining of paraffin embedded Human Placenta, with no primary antibody.




Immunohistochemistry-Paraffin: u-Plasminogen Activator/Urokinase Antibody [NB100-2419] - Staining of paraffin embedded Human Prostate. Antibody at 8 ug/mL. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



Immunohistochemistry-Paraffin: u-Plasminogen Activator/Urokinase Antibody [NB100-2419] - Negative Control showing staining of paraffin embedded Human Prostate, with no primary antibody.



Immunohistochemistry-Paraffin: u-Plasminogen Activator/Urokinase Antibody [NB100-2419] - Staining of paraffin embedded Human Placenta. Antibody at 8 ug/mL. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



Publications

Morgan H, Hill PA. Human breast cancer cell-mediated bone collagen degradation requires plasminogen activation and matrix metalloproteinase activity. *Cancer Cell Int* 2005-02-08 [PMID: 15701164]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-2419

NB800-PC6	293 Whole Cell Lysate
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-2419

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



