# **Product Datasheet**

# BCL11B Antibody NB100-2600

Unit Size: 0.1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 7

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-2600

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB100-2600



## NB100-2600


**BCL11B** Antibody

Unit Size 0.1 ml Concentration 0.25 mg/ml Storage Store at 4C. Do not freeze. Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer TBS and 0.1% BSA  Product Description Host Rabbit Gene ID 64919 Gene Symbol BCL11B Species Human, Mouse, Rat Reactivity Notes Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%). Immunogen Tecopicalized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details Applications Immunocytochemistry/ Immunofluorescence, Immunohistochemistry/ Paraffin 1:100-1:500, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Prozen, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemi	BOET 1B / Millibody	
Concentration  Storage  Store at 4C. Do not freeze.  Clonality  Polyclonal  Preservative  0.09% Sodium Azide  Isotype  IgG  Purity  Immunogen affinity purified  Buffer  TBS and 0.1% BSA  Product Description  Host  Gene ID  64919  Gene Symbol  BCL11B  Species  Human, Mouse, Rat  Reactivity Notes  Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%).  Immunogen  The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLLI/ymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details  Applications  Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin  Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin  Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin  Recommended Dilutions  Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity re	Product Information	
Storage Store at 4C. Do not freeze.  Clonality Polyclonal  Preservative 0.09% Sodium Azide  Isotype IgG  Purity Immunogen affinity purified  Buffer TBS and 0.1% BSA  Product Description  Host Rabbit Gene ID 64919  Gene Symbol BCL11B  Species Human, Mouse, Rat  Reactivity Notes Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%).  Immunogen The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLLI/ymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details  Applications Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin  Recommended Dilutions Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155	Unit Size	0.1 ml
Clonality Polyclonal Preservative 0.09% Sodium Azide Isotype IgG Purity Immunogen affinity purified Buffer TBS and 0.1% BSA  Product Description Host Rabbit Gene ID 64919 Gene Symbol BCL11B Species Human, Mouse, Rat Reactivity Notes Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%). Immunogen The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details Applications Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Recommended Dilutions Immunocytochemistry 1:100-1:500, Immunocytochemistry/ Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen Application Notes Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Concentration	0.25 mg/ml
Preservative  Isotype  IgG  Purity  Immunogen affinity purified  Buffer  TBS and 0.1% BSA  Product Description  Host  Rabbit  Gene ID  Gene Symbol  BCL11B  Species  Human, Mouse, Rat  Reactivity Notes  Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%).  Immunogen  The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details  Applications  Immunocytochemistry/ Immunofluorescence, Immunohistochemistry/ Immunohistochemistry-Paraffin  Recommended Dilutions  Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes  Epitope exposure is recommended to enhance staining, ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Storage	Store at 4C. Do not freeze.
Isotype	Clonality	Polyclonal
Purity Immunogen affinity purified  Buffer TBS and 0.1% BSA  Product Description  Host Rabbit  Gene ID 64919  Gene Symbol BCL11B  Species Human, Mouse, Rat  Reactivity Notes Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%).  Immunogen The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details  Applications Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin  Recommended Dilutions Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunofluorescence, Immunohistochemistry-Prozen  Application Notes Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Preservative	0.09% Sodium Azide
Product Description  Host Rabbit  Gene ID 64919  Gene Symbol BCL11B  Species Human, Mouse, Rat  Reactivity Notes Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%).  Immunogen The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details  Applications Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin  Recommended Dilutions Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Isotype	IgG
Product Description  Host Rabbit  Gene ID 64919  Gene Symbol BCL11B  Species Human, Mouse, Rat  Reactivity Notes Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%).  Immunogen The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details  Applications Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin  Recommended Dilutions Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Purity	Immunogen affinity purified
Host Rabbit Gene ID 64919 Gene Symbol BCL11B Species Human, Mouse, Rat Reactivity Notes Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%). Immunogen The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details Applications Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Prozen Application Notes Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Buffer	TBS and 0.1% BSA
Gene Symbol  Gene Symbol  BCL11B  Species  Human, Mouse, Rat  Reactivity Notes  Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%).  Immunogen  The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details  Applications  Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin  Recommended Dilutions  Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes  Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Product Description	
Gene Symbol  Species  Human, Mouse, Rat  Reactivity Notes  Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%).  Immunogen  The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details  Applications  Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin  Recommended Dilutions  Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes  Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Host	Rabbit
Reactivity Notes   Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%).	Gene ID	64919
Reactivity Notes  Rat reactivity reported in scientific literature (PMID: 26465886). Mouse reactivity reported from a verified customer review. Gorilla, X.tropicalis (100%).  The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details  Applications  Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin  Recommended Dilutions  Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes  Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Gene Symbol	BCL11B
reported from a verified customer review. Gorilla, X.tropicalis (100%).  Immunogen  The immunogen recognized by this antibody maps to a region between residue 850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details  Applications  Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin  Recommended Dilutions  Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes  Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Species	Human, Mouse, Rat
850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using the numbering given in Swiss-Prot entry Q9C0K0 (GeneID 64919).  Product Application Details  Applications  Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin  Recommended Dilutions  Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes  Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Reactivity Notes	
Applications  Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin  Recommended Dilutions  Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes  Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Immunogen	850 and the C-terminus (residue 894) of human B-cell CLL/lymphoma 11B using
Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin  Recommended Dilutions  Immunohistochemistry 1:100-1:500, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes  Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Product Application Details	
Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500, Immunohistochemistry-Frozen  Application Notes  Epitope exposure is recommended to enhance staining. ICC/IF reactivity reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Applications	
reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from	Recommended Dilutions	Immunofluorescence, Immunohistochemistry-Paraffin 1:100-1:500,
	Application Notes	reported in scientific literature (PMID: 24155294). IHC-F reactivity reported in scientific literature (PMID: 24155294). BCL11B antibody validated for IHC-P from




### **Images**


Immunohistochemistry-Paraffin: BCL11B Antibody [NB100-2600] - Section of human skin tumor (higher magnification). Antibody: Affinity purified rabbit anti-Bcl11b used at a dilution of 1:500. Detection: DAB staining using anti-Rabbit IHC antibody at a dilution of 1:100.




Immunohistochemistry-Paraffin: BCL11B Antibody [NB100-2600] - P7 brain mouse section was stained with anti-BCL11B antibody. Image from verified customer review.



Immunohistochemistry: BCL11B Antibody [NB100-2600] - Immunohistochemistry and FISH of AML cases. Immunohistochemistry analysis of BCL11B-expressing AML samples #59810, #57863 and #13 carrying t(2;14)(q22.3;q32.3), t(7;14)(q21q32) and no 14q32 alteration, respectively. BCL11B expression was detected in samples regardless of the presence of the 14q32 alterations. The expression was limited to the nucleus and the percentage of positive neoplastic cells was always >= 50%. Image collected and cropped by CiteAb from the following publication (https://www.mdpi.com/2072-6694/11/12/1951), licensed under a CC-BY license.



Immunohistochemistry-Paraffin: BCL11B Antibody [NB100-2600] - Section of human skin tumor. Antibody: Affinity purified rabbit anti-Bcl11b used at a dilution of 1:500. Detection: DAB staining using anti-Rabbit IHC antibody at a dilution of 1:100.



#### **Publications**

Wang Y, Zhang F, Gao N et al. Differential expression of BCL11b and CDKN2A in CD30-positive peripheral T cell lymphoma: Retrospective study Medicine 2023-11-17 [PMID: 37986346] (IHC-P, Human)

Bakalar D, O'Reilly JJ, Lacaille H et al. Lack of placental neurosteroid alters cortical development and female somatosensory function Front Endocrinol (Lausanne) 2022-10-13 [PMID: 36313771]

Traunm□ller L, Schulz J, Ortiz R et al. A cell-type-specific alternative splicing regulator shapes synapse properties in a trans-synaptic manner Cell reports 2023-02-28 [PMID: 36862556] (IHC, Mouse)

Padella A, Simonetti G, Paciello G, et al. Novel and Rare Fusion Transcripts Involving Transcription Factors and Tumor Suppressor Genes in Acute Myeloid Leukemia Cancers (Basel) 2019-12-05 [PMID: 31817495] (IF/IHC, Human)

Digilio L, Yap CC, Winckler B. Ctip2-, Satb2-, Prox1-, and GAD65-Expressing Neurons in Rat Cultures: Preponderance of Single- and Double-Positive Cells, and Cell Type-Specific Expression of Neuron-Specific Gene Family Members, Nsg-1 (NEEP21) and Nsg-2 (P19). PLoS One 2015-01-01 [PMID: 26465886] (Rat)

Carpentier PA, Haditsch U, Braun AE et al. Stereotypical alterations in cortical patterning are associated with maternal illness-induced placental dysfunction. J Neurosci. 2013-10-23 [PMID: 24155294] (IHC-Fr, ICC/IF, Mouse)

Beguin S, Crepel V, Aniksztejn L, Becq H, Pelosi B, Pallesi-Pocachard E, Bouamrane L, Pasqualetti M, Kitamura K, Cardoso C, Represa A. An Epilepsy-Related ARX Polyalanine Expansion Modifies Glutamatergic Neurons Excitability and Morphology Without Affecting GABAergic Neurons Development. Cereb Cortex. 2012-05-24 [PMID: 22628459] (IF/IHC, Mouse)





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

## **Products Related to NB100-2600**

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-33549PEP BCL11B Recombinant Protein Antigen

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-2600

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

