

Product Datasheet

PBR Antibody NB100-41398

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 13

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-41398

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-41398



NB100-41398

PBR Antibody

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA
Target Molecular Weight	18.8 kDa

Product Description	
Description	Novus Biologicals Goat PBR Antibody (NB100-41398) is a polyclonal antibody validated for use in IHC, WB, ELISA, Flow, ICC/IF and IP. Anti-PBR Antibody: Cited in 11 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	706
Gene Symbol	TSPO
Species	Human, Mouse, Rat
Reactivity Notes	Use in Mouse reported in scientific literature (PMID:33725191).
Specificity/Sensitivity	This antibody is expected to recognise one isoform (NP_000705.2) only.
Immunogen	Peptide with sequence C-RDNHGWRGGRRLE corresponding to C-Terminus according to NP_000705.2.


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Peptide ELISA
Recommended Dilutions	Western Blot 0.1-0.3 ug/ml, Flow Cytometry 10 ug/mL, Immunohistochemistry 2-4 ug/ml, Immunocytochemistry/ Immunofluorescence 10 ug/mL, Immunohistochemistry-Paraffin 2-4 ug/ml, Peptide ELISA detection limit 1:128000

Images


Western Blot: PBR Antibody [NB100-41398] - Staining of A431 cell lysate.(35 ug protein in RIPA buffer). Antibody at 0.3 ug/mL. Detected by chemiluminescence.




Immunocytochemistry/Immunofluorescence: PBR Antibody [NB100-41398] - Immunofluorescence analysis of paraformaldehyde fixed MCF7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL).




Immunohistochemistry: PBR Antibody [NB100-41398] - Correlative TSPO immunohistochemistry in a primary human grade III astrocytoma xenotransplant, Tumor + White Matter Tract (40X). Image collected and cropped by CiteAb from the following publication (www.journals.plos.org/plosone/article?id=10.1371/journal.pone.0141659) licensed under a CC-BY license.




Flow Cytometry: PBR Antibody [NB100-41398] - Flow cytometric analysis of paraformaldehyde fixed MCF7 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (1 ug/mL). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.




Immunocytochemistry/Immunofluorescence: PBR Antibody [NB100-41398] - Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative cControl: Unimmunized goat IgG (10 ug/mL) followed by Alexa Fluor 488 secondary antibody.




Immunohistochemistry-Paraffin: PBR Antibody [NB100-41398] - Staining of paraffin embedded Human Colon. Antibody at 2.5 ug/mL. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.




Immunohistochemistry-Paraffin: PBR Antibody [NB100-41398] - Staining of paraffin embedded Human Kidney. Antibody at 5 ug/mL. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.




Immunohistochemistry: PBR Antibody [NB100-41398] - Human tumor microarrays (TMAs). Pilocytic astrocytoma TSPO immunohistochemistry (A) and Glioblastoma multiforme TSPO immunohistochemistry (B). Image collected and cropped by CiteAb from the following publication (www.journals.plos.org/plosone/article?id=10.1371/journal.pone.0141659) licensed under a CC-BY license.




Immunohistochemistry: PBR Antibody [NB100-41398] - [18F]PBR06 selectivity in primary human grade IV glioblastoma xenotransplant. (A) Dynamic PET image (axial) pre-infusion of cold analog, with [18F]PBR06 uptake primarily confined to the tumor. (B) Dynamic PET image (axial) post-infusion, showing nearly total displacement of [18F]PBR06. (C) Time-activity curves of injected [18F]PBR06 in tumor (green) & contralateral brain (blue). (D) Correlative TSPO immunohistochemistry, Tumor + White Matter Tract (40X). Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0141659>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry-Paraffin: PBR Antibody [NB100-41398] - Human tumor microarrays (TMAs).Pilocytic astrocytoma TSPO immunohistochemistry (A) & TSPO immunohistochemical scoring (C). Glioblastoma multiforme TSPO immunohistochemistry (B) & TSPO immunohistochemical scoring (D). Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0141659>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry-Paraffin: PBR Antibody [NB100-41398] - Human tumor microarrays (TMAs).Pilocytic astrocytoma TSPO immunohistochemistry (A) & TSPO immunohistochemical scoring (C). Glioblastoma multiforme TSPO immunohistochemistry (B) & TSPO immunohistochemical scoring (D). Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0141659>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry-Paraffin: PBR Antibody [NB100-41398] - Primary human grade IV glioblastoma xenotransplant.(A) Dynamic PET image (coronal) of advanced tumor in the right hemisphere of the brain with [18F]PBR06 uptake confined primarily to the tumor. Arrows indicate tumor & infiltration into left hemisphere. (B) Time-activity curves of injected [18F]PBR06 in tumor (green) & contralateral brain (blue) [30]. (C) Correlative TSPO immunohistochemistry, Tumor + White Matter Tract (40X). Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0141659>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Immunohistochemistry-Paraffin: PBR Antibody [NB100-41398] - Primary human grade III astrocytoma xenotransplant.(A) T2-weighted MRI (coronal) visualizes advanced tumor in the right hemisphere of the brain. (B) Correlative dynamic PET image (coronal) of same advanced tumor in the right hemisphere of the brain, with [18F]PBR06 uptake primarily confined to the tumor & co-localizing with tumor tissue visualized by MRI. Top arrows indicate tumor & infiltration into left hemisphere. (C) Fused MRI (A) PET (B) image. (D) Time-activity curves of injected [18F]PBR06 in tumor (green) & contralateral brain (blue). Correlative vimentin immunohistochemistry: (E) Gross; (F) Tumor + White Matter Tract (40X); (G) Tumor (40X). (H) Correlative dynamic PET image, axial view along yellow line in (B); arrows indicate tumor & infiltration into left hemisphere. Correlative TSPO immunohistochemistry: (I) Gross; (J) Tumor + White Matter Tract (40X); (K) Tumor (40X). (L) Dynamic PET image (axial) of control cohort. Correlative CD68 immunohistochemistry: (M) Gross; (N) Tumor + White Matter Tract (40X); (O) Tumor (40X). Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0141659>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunohistochemistry-Paraffin: PBR Antibody [NB100-41398] - Primary human grade III astrocytoma xenotransplant. (A) T2-weighted MRI (coronal) visualizes advanced tumor in the right hemisphere of the brain. (B) Correlative dynamic PET image (coronal) of same advanced tumor in the right hemisphere of the brain, with [18F]PBR06 uptake primarily confined to the tumor & co-localizing with tumor tissue visualized by MRI. Top arrows indicate tumor & infiltration into left hemisphere. (C) Fused MRI (A) PET (B) image. (D) Time-activity curves of injected [18F]PBR06 in tumor (green) & contralateral brain (blue). Correlative vimentin immunohistochemistry: (E) Gross; (F) Tumor + White Matter Tract (40X); (G) Tumor (40X). (H) Correlative dynamic PET image, axial view along yellow line in (B); arrows indicate tumor & infiltration into left hemisphere. Correlative TSPO immunohistochemistry: (I) Gross; (J) Tumor + White Matter Tract (40X); (K) Tumor (40X). (L) Dynamic PET image (axial) of control cohort. Correlative CD68 immunohistochemistry: (M) Gross; (N) Tumor + White Matter Tract (40X); (O) Tumor (40X). Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0141659>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Immunohistochemistry: PBR Antibody [NB100-41398] - Primary human grade III astrocytoma xenotransplant. (A) T2-weighted MRI (coronal) visualizes advanced tumor in the right hemisphere of the brain. (B) Correlative dynamic PET image (coronal) of same advanced tumor in the right hemisphere of the brain, with [18F]PBR06 uptake primarily confined to the tumor & co-localizing with tumor tissue visualized by MRI. Top arrows indicate tumor & infiltration into left hemisphere. (C) Fused MRI (A) PET (B) image. (D) Time-activity curves of injected [18F]PBR06 in tumor (green) & contralateral brain (blue). Correlative vimentin immunohistochemistry: (E) Gross; (F) Tumor + White Matter Tract (40X); (G) Tumor (40X). (H) Correlative dynamic PET image, axial view along yellow line in (B); arrows indicate tumor & infiltration into left hemisphere. Correlative TSPO immunohistochemistry: (I) Gross; (J) Tumor + White Matter Tract (40X); (K) Tumor (40X). (L) Dynamic PET image (axial) of control cohort. Correlative CD68 immunohistochemistry: (M) Gross; (N) Tumor + White Matter Tract (40X); (O) Tumor (40X). Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0141659>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

- Cohen AS, Li J, Hight MR et al. TSPO-targeted PET and Optical Probes for the Detection and Localization of Premalignant and Malignant Pancreatic Lesions Clinical Cancer Research 2020-11-15 [PMID: 32933996]
- Nutma E, Fancy N, Marzin M et al. Translocator protein is a marker of activated microglia in rodent models but not human neurodegenerative diseases Research Square 2022-03-23
- Nutma E, Gebro E, Marzin MC et al. Activated microglia do not increase 18 kDa translocator protein (TSPO) expression in the multiple sclerosis brain Glia 2021-06-19 [PMID: 34145928]
- Chang CW, Chiu CH, Lin MH et al. GMP-compliant fully automated radiosynthesis of [18F]FEPPA for PET/MRI imaging of regional brain TSPO expression EJNMMI research 2021-03-16 [PMID: 33725191] (WB, ICC/IF, Mouse)
- Robb J Characterisation of immunometabolic responses in astrocytes Thesis 2020-01-01 (IP)
- Hieu Tran V, Park H, Park J et al. Synthesis and evaluation of novel potent TSPO PET ligands with 2-phenylpyrazolo [1,5-a]pyrimidin-3-yl acetamide Bioorg. Med. Chem. 2019-07-19 [PMID: 31353076] (IF/IHC, Rat, Human)
- Buck JR, McKinley ET, Fu A et al. Preclinical TSPO Ligand PET to Visualize Human Glioma Xenotransplants: A Preliminary Study. PLoS ONE. 2015-10-31 [PMID: 26517124] (IHC-P, Human)
- Papadopoulos V, Baraldi M, Guilarte TR et al. Translocator protein (18kDa): new nomenclature for the peripheral-type benzodiazepine receptor based on its structure and molecular function. Trends Pharmacol Sci 2006-08-01 [PMID: 16822554]
- Zhou T, Frabutt DA, Moremen KW, Zheng YH et al. ERMAnI is required for HIV-1 envelope glycoprotein degradation via endoplasmic reticulum-associated protein degradation pathway J. Biol. Chem. 2015-07-23 [PMID: 26205822] (WB, Human)
- Zhou T, Dang Y, Zheng YH. The Mitochondrial Translocator Protein, TSPO, Inhibits HIV-1 Envelope Glycoprotein Biosynthesis via the Endoplasmic Reticulum-Associated Protein Degradation Pathway. J. Virol. 2014-03-01 [PMID: 24403586] (WB, Human)
- Wu X, Gallo KA. The 18-kDa Translocator Protein (TSPO) Disrupts Mammary Epithelial Morphogenesis and Promotes Breast Cancer Cell Migration. PLoS One. 2013-08-14 [PMID: 23967175] (WB, ICC/IF, Human)
- Joo, HK et al. Peripheral benzodiazepine receptor regulates vascular endothelial activations via suppression of the voltage-dependent anion channel-1 FEBS Lett 2012-01-01 [PMID: 22616995] (WB, Human)
- More publications at <http://www.novusbio.com/NB100-41398>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-41398

NBL1-17384	PBR Overexpression Lysate
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-41398

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



