Product Datasheet

PCNA Antibody - BSA Free NB100-456

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com

G

technical@novusbio.com

Publications: 7

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-456

Updated 2/21/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB100-456



NB100-456

PCNA Antibody - BSA Free

,	
Product Information	
Unit Size	100 ul
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)
Product Description	
Host	Rabbit
Gene ID	5111
Gene Symbol	PCNA
Species	Human, Mouse
Marker	Proliferation Marker
Immunogen	The immunogen this antibody was made to, maps to a region between residues 75-125 of human proliferating cell nuclear antigen using the SwissProt entry P12004 (GeneID 5111).
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry- Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000-1:10000, Flow Cytometry 1:10-1:1000, Immunohistochemistry 1:2000 - 1:10000, Immunoprecipitation 2-5 ug/mg lysate, Immunohistochemistry-Paraffin 1:2000 - 1:10000
Application Notes	Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.


Images


Western Blot: PCNA Antibody [NB100-456] - Detection of Human and Mouse PCNA by Western Blot. Samples: Whole cell lysate (50 ug) from 293T and mouse NIH3T3 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-PCNA antibody NB100-456 used for WB at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.

kDa	
kDa 250-	
130	
70-	
51-	
38-	
28-	- PCNA
19-	
16-	
293T 3T3	



Immunohistochemistry: PCNA Antibody [NB100-456] - Sample: FFPE section of human stomach carcinoma. Antibody: Affinity purified rabbit anti- PCNA used at a dilution of 1:10,000 (0.1ug/ml). Detection: DAB






www.novusbio.com



Immunoprecipitation: PCNA Antibody [NB100-456] - Detection of human PCNA by western blot and immunoprecipitation. Samples: A. Whole cell lysate (WCL) from A-431 cells. B. WCL (500 ug for IP; 30 ug for WCL lane) from MDA-MB-468 cells. Antibodies: A. Affinity purified rabbit anti-PCNA antibody NB100-456 used at the indicated concentrations for WB. B. PCNA was immunoprecipitated using another a mouse monoclonal antibody to PCNA (mAb). Control mock IP was performed using normal mouse IgG (M-IgG) and normal rabbit IgG (R-IgG). immunoprecipitatesd PCNA was blotted using NB100-456 (1 ug/ml). Detection: Chemiluminescence with an exposure time of 2 minutes (A) or 5 seconds (B).



Publications

Chakrabarti J, Holokai L, Syu L et al. Hedgehog signaling induces PD-L1 expression and tumor cell proliferation in gastric cancer. Oncotarget 2018-12-21 [PMID: 30647844]

Yang TY, Wu YL, Yu MH et al. Mulberry Leaf and Neochlorogenic Acid Alleviates Glucolipotoxicity-Induced Oxidative Stress and Inhibits Proliferation/Migration via Downregulating Ras and FAK Signaling Pathway in Vascular Smooth Muscle Cell Nutrients 2022-07-22 [PMID: 35893859] (WB, Rat)

Koguchi M, Nakahara Y, Ito H et al. BMP4 induces asymmetric cell division in human glioma stem-like cells Oncol Lett 2020-02-01 [PMID: 31966054] (WB, Human)

Lo YH, Ho PC, Wang SC et al. Epidermal growth factor receptor protects proliferating cell nuclear antigen from cullin 4A protein-mediated proteolysis. J Biol Chem 2012-08-01 [PMID: 22692198]

Zhu Y, Wu Z, Cardoso MC et al. Processing of lagging-strand intermediates in vitro by herpes simplex virus type 1 DNA polymerase. J Virol 2010-08-01 [PMID: 20444887]

Ramos-Nino ME, Blumen SR, Sabo-Attwood T et al. HGF mediates cell proliferation of human mesothelioma cells through a PI3K/MEK5/Fra-1 pathway. Am J Respir Cell Mol Biol 2008-02-01 [PMID: 17872495]

Williams SA, Longerich S, Sung P et al. The E3 ubiquitin ligase RAD18 regulates ubiquitylation and chromatin loading of FANCD2 and FANCI. Blood 2011-05-01 [PMID: 21355096]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NB100-456

NBL1-14184	PCNA Overexpression Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-456

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

