

Product Datasheet

USP7 Antibody - BSA Free NB100-513

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 30


Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-513

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-513



NB100-513

USP7 Antibody - BSA Free

Product Information	
Unit Size	100 ul
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)
Target Molecular Weight	128 kDa


Product Description	
Host	Rabbit
Gene ID	7874
Gene Symbol	USP7
Species	Human, Mouse
Immunogen	The immunogen recognized by this antibody maps to a region between residues 1 and 50 of human Ubiquitin Specific Peptidase 7 using the numbering given in entry NP_003461.1 (GeneID 7874).

Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:10000-1:25000, Immunohistochemistry 1:1000 - 1:5000, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation 2-10 ug/ml of lysate, Immunohistochemistry-Paraffin 1:1000-1:5000
Application Notes	Many tissues will not require epitope retrieval. However to boost weak staining, epitope retrieval with citrate buffer at pH 6.0 is recommended for FFPE tissue sections. USP7 antibody validated for ICC/IF from a verified customer review.




Images


Western Blot: USP7 Antibody [NB100-513] - Samples: Whole cell lysate (10 ug) from HEK293T, RKO, Hep-G2, Jurkat, HeLa, TCMK-1, and NIH 3T3 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-USP7 antibody used for WB at 0.04 ug/ml. Detection: Chemiluminescence with an exposure time of 10 seconds




Immunohistochemistry-Paraffin: USP7 Antibody [NB100-513] - Sample: FFPE section of human ovarian carcinoma. Antibody: Affinity purified rabbit anti-USP7 used at 1:5,000 (0.2ug/ml). Detection: DAB



Immunohistochemistry-Paraffin: USP7 Antibody [NB100-513] - Sample: FFPE section of mouse renal cell carcinoma. Antibody: Affinity purified rabbit anti-USP7 used at 1:5,000 (0.2ug/ml). Detection: DAB



Immunoprecipitation: USP7 Antibody [NB100-513] - Samples: Whole cell lysate (10 ug) from HEK293T, RKO, Hep-G2, Jurkat, HeLa, TCMK-1, and NIH 3T3 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-USP7 antibody used for WB at 0.04 ug/ml. Detection: Chemiluminescence with an exposure time of 10 seconds



Publications

Gao A, Zhang M, Zhu SQ et al. DNA polymerase iota promotes EMT and metastasis of esophageal squamous cell carcinoma by interacting with USP7 to stabilize HIF-1? *Cell Death Dis* 2024-02-24 [PMID: 38402183]

Liu J, Zhou T, Dong X et al. De-ubiquitination of SAMHD1 by USP7 promotes DNA damage repair to overcome oncogenic stress and affect chemotherapy sensitivity *Oncogene* 2023-04-20 [PMID: 37081042] (IHC, Human)

Liu J, Zhou T, dong x Et al. De-ubiquitination of SAMHD1 by USP7 overcomes oncogenic stress and contributes to chemotherapy insensitivity *Research Square* 2021-08-20 (IF/IHC, Human)

Nicklas S, Hillje AL, Okawa S et al. A complex of the ubiquitin ligase TRIM32 and the deubiquitinase USP7 balances the level of c-Myc ubiquitination and thereby determines neural stem cell fate specification *Cell Death Differ.* 2018 Jun 13 [PMID: 29899379] (IHC, Mouse)

Nicklas S, Hillje AL, Okawa S et al. A complex of the ubiquitin ligase TRIM32 and the deubiquitinase USP7 balances the level of c-Myc ubiquitination and thereby determines neural stem cell fate specification *Cell Death Differ.* 2018 Jun 13 [PMID: 29899379] (IHC, Mouse)

Cui H, Guo M, Xu D et al. The stress-responsive gene ATF3 regulates the histone acetyltransferase Tip60 *Nat Commun* 2015-04-13 [PMID: 25865756] (IP, WB, Human)

de Lange J, Teunisse AF, Vries MV et al. High levels of Hdmx promote cell growth in a subset of uveal melanomas. *Am J Cancer Res* 2012-01-01 [PMID: 22957303]

Fei J, Chen J. KIAA1530 protein is recruited by Cockayne syndrome complementation group protein A (CSA) to participate in transcription-coupled repair (TCR). *J Biol Chem* 2012-10-01 [PMID: 22902626]

Cano F, Bye H, Duncan LM et al. The RNA-binding E3 ubiquitin ligase MEX-3C links ubiquitination with MHC-I mRNA degradation. *EMBO J* 2012-08-01 [PMID: 22863774]

Dekker P, Gunn D, McBryan T et al. Microarray-based identification of age-dependent differences in gene expression of human dermal fibroblasts. *Mech Ageing Dev* 2012-07-01 [PMID: 22721680]

Lenos K, Grawenda AM, Lodder K et al. Alternate splicing of the p53 inhibitor HDMX offers a superior prognostic biomarker than p53 mutation in human cancer. *Cancer Res* 2012-08-01 [PMID: 22700878]

Schwertman P, Lagarou A, Dekkers DH et al. UV-sensitive syndrome protein UVSSA recruits USP7 to regulate transcription-coupled repair. *Nat Genet* 2012-05-01 [PMID: 22466611]

More publications at <http://www.novusbio.com/NB100-513>





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-513

NBL1-17671	USP7 Overexpression Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-513

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

