# **Product Datasheet**

# Caspase-6 Antibody NB100-56114

Unit Size: 0.05 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

**Publications: 4** 

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-56114

Updated 4/21/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB100-56114



# NB100-56114

Caspase-6 Antibody

Product Information	
0.05 ml	
This product is unpurified. The exact concentration of antibody is not quantifiable.	
Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.	
Polyclonal	
0.05% Sodium Azide	
IgG	
Unpurified	
50 ul neat whole antisera	
Product Description	
Rabbit	
CASP6	
Human, Mouse, Rat, Canine	
Cross reacts with Dog, Gerbil, Human, Mouse and Rat. Not yet tested in other species.	
This polyclonal antisera recognizes the large and small subunits of active/cleaved caspase-6. Whereas the antisera has a strong preference for active/cleaved caspase-6, in some cell or tissue systems or techniques the antisera may also recognize the ~34 kDa proform of caspase-6 as well as intermediate caspase-6 cleavage fragments.	
Recombinant catalytically active human Caspase-6 protein was used as immunogen.	
Product Application Details	
Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry- Paraffin, Immunoprecipitation	
Western Blot 1:1000-1:2000, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation 1:50-1:200, Immunohistochemistry- Paraffin 1:1000-1:5000, Immunohistochemistry-Frozen	
Immunohistochemistry (frozen): Users should optimize according to model and immunodetection system used (secondary reagents).	

www.novusbio.com




#### Images

Western Blot: Caspase-6 Antibody [NB100-56114] - Analysis of Caspase-6 using this antibody. Recombinant catalytically active Caspase-6 was western blotted with Active/Cleaved Caspase-6 antibody. The antibody detected both the large and small subunits of Caspase-6.

Immunohistochemistry-Paraffin: Caspase-6 Antibody [NB100-56114] -Formalin-fixed paraffin-embedded tissue sections of human gliomas stained for Active/Cleaved Caspase-6 expression using Caspase-6 antibody at 1:2000. Gemistocytoma (low grade tumor) at A) low and B) high magnification. Two examples of anaplastic, high grade (III) infiltrating gliomas (C and D). Increased active/cleaved Caspase-6 may be seen in high grade, compared to low grade tumors. Hematoxylineosin counterstain.

Immunohistochemistry: Caspase-6 Antibody [NB100-56114] - Formalinfixed paraffin-embedded tissue section of dog cerebellum stained for Caspase-6 expression at 1:2000. Hematoxylin-eosin counterstain.





#### **Publications**

Berta T, Park CK, Xu ZZ et al. Extracellular caspase-6 drives murine inflammatory pain via microglial TNF-alpha secretion. J. Clin. Invest. 2014-03-03 [PMID: 24531553] (WB, Mouse)

#### Details:

WB (mouse spinal cord tissue, cerebral fluid and recombinant Caspase-6), Figs 2C-F, S1. The Caspase-6 pAb was used at 1:2000. The specificity of the antibody was validated with recombinant caspase-6 by WB (Fig 2E). Procaspase-6 was detected at ~37kDa, an

Saggar JK, Tannock IF. Activity of the hypoxia-activated pro-drug TH-302 in hypoxic and perivascular regions of solid tumors and its potential to enhance therapeutic effects of chemotherapy. Int J Cancer. 2013-11-08 [PMID: 24338277] (IHC-Fr, ICC/IF, Human)

Krajewska M, Rosenthal RE, Mikolajczyk J et al. Early processing of Bid and caspase-6, -8, -10, -14 in the canine brain during cardiac arrest and resuscitation. Exp Neurol. 2004-10-01 [PMID: 15380478] (IHC-P, WB, Rat, Canine, Mouse)

#### Details:

Antibodies cited: 1. Caspase-9 (Active/Cleaved), IMG-5705: WB: Fig 1A (recombinant human caspase-9), Fig 4A (recombinant human caspase-9, rat brain), Fig 4B-F (isolated mitochondria from rat liver or heart). Fig 5 (PC12 cells) IHC (P): Fig 2d-k (rat kidne

Saggar JK, Fung AS, Patel KJ, Tannock IFet al. Use of molecular biomarkers to quantify the spatial distribution of effects of anticancer drugs in solid tumors. Mol Cancer Ther 2013-04-01 [PMID: 23348047] (ICC/IF, IF/IHC, Human)





# Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

# **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

# **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

## Products Related to NB100-56114

NBP3-11853	Jurkat Staurosporine Treated / Untreated Cell Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-56114

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

