

Product Datasheet

DGAT1 Antibody NB100-57086

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 10

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-57086

Updated 9/9/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-57086



NB100-57086**DGAT1 Antibody**

Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA
Target Molecular Weight	55.3 kDa

Product Description	
Description	Novus Biologicals Goat DGAT1 Antibody (NB100-57086) is a polyclonal antibody validated for use in IHC, WB, ELISA, Flow and ICC/IF. Anti-DGAT1 Antibody: Cited in 10 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Goat
Gene ID	8694
Gene Symbol	DGAT1
Species	Human, Mouse
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 25714675).
Immunogen	Peptide with sequence C-QNSMKPFKDM DYS corresponding to internal region according to NP_036211.1.

Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Peptide ELISA
Recommended Dilutions	Western Blot 0.01 - 0.1 ug/mL, Flow Cytometry 10 ug/mL, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence 10 ug/mL, Immunohistochemistry-Paraffin 3.75 ug/mL, Peptide ELISA Detection limit 1:64000
Application Notes	WB: Approx. 55 kDa band observed in human liver lysates (calculated MW of 55.3 kDa band according to NP_036211.1).




Images


Western Blot: DGAT1 Antibody [NB100-57086] - Staining of Mouse Liver lysate (35 ug protein in RIPA buffer). Antibody at 0.03 ug/mL. Detected by chemiluminescence.

250kDa
150kDa
100kDa
75kDa
50kDa
37kDa
25kDa
20kDa
15kDa


Immunocytochemistry/Immunofluorescence: DGAT1 Antibody [NB100-57086] - Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL), showing endoplasmic reticulum and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (2 ug/mL).



Immunohistochemistry-Paraffin: DGAT1 Antibody [NB100-57086] - Staining of paraffin embedded Human Colon. Antibody at 3.75 ug/mL. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Flow Cytometry: DGAT1 Antibody [NB100-57086] - Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 ug/mL) followed by Alexa Fluor 488 secondary antibody (1 ug/mL). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



Publications

Larsen Steen, Vigelso Andreas, Dandanell Sune et al. Simvastatin-Induced Insulin Resistance May Be Linked to Decreased Lipid Uptake and Lipid Synthesis in Human Skeletal Muscle: the LIFESTAT Study. J Diabetes Res. 2018-09-12 [PMID: 30276217]

Melissa L. Erickson, Zachary W. Patinkin, Allison M. Duensing, Dana Dabelea, Leanne M. Redman, Kristen E. Boyle Maternal metabolic health drives mesenchymal stem cell metabolism and infant fat mass at birth JCI Insight 2021-07-08 [PMID: 34061777]

Timothy M. Moore, Zhenqi Zhou, Alexander R. Strumwasser, Whitaker Cohn, Amanda J. Lin, Kevin Cory, Kate Whitney, Theodore Ho, Timothy Ho, Joseph L. Lee, Daniel H. Rucker, Austin N. Hoang, Kevin Widjaja, Aaron D. Abrishami, Sarada Charugundla, Linsey Stiles, Julian P. Whitelegge, Lorraine P. Turcotte, Jonathan Wanagat, Andrea L. Hevener Age-induced mitochondrial DNA point mutations are inadequate to alter metabolic homeostasis in response to nutrient challenge Aging Cell 2020-10-13 [PMID: 33049094]

Chaves AB, Zheng D, Johnson JA et al. Infant Mesenchymal Stem Cell Insulin Action is Associated With Maternal Plasma Free Fatty Acids, Independent of Obesity Status: The Healthy Start Study Diabetes 2022-05-27 [PMID: 35621990]

Bagnato C, Igal RA. Overexpression of diacylglycerol acyltransferase-1 reduces phospholipid synthesis, proliferation, and invasiveness in simian virus 40-transformed human lung fibroblasts. J Biol Chem 2003-12-26 [PMID: 14557275]

Borsting Jordy A, Kraakman MJ, Gardner T et al. Analysis of the liver lipidome reveals insights into the protective effect of exercise on high fat diet induced hepatosteatosis in mice Am. J. Physiol. Endocrinol. Metab. 2015-02-24 [PMID: 25714675] (WB, Mouse)

Seyer A, Cantiello M, Bertrand-Michel J et al. Lipidomic and Spatio-Temporal Imaging of Fat by Mass Spectrometry in Mice Duodenum during Lipid Digestion PLoS One 2013-01-01 [PMID: 23560035] (WB, Mouse)

van Diepen JA, Stienstra R, Vroegrijk IO et al. Caspase-1 deficiency in mice reduces intestinal triglyceride absorption and hepatic triglyceride secretion. J Lipid Res 2013-02-01 [PMID: 23160218]

Amati F, Dube JJ, Alvarez-Carnero E et al. Skeletal muscle triglycerides, diacylglycerols, and ceramides in insulin resistance: another paradox in endurance-trained athletes? Diabetes 2011-10-01 [PMID: 21873552] (WB, Human)

Alsted TJ, Nybo L, Schweiger M et al. Adipose triglyceride lipase in human skeletal muscle is upregulated by exercise training. Am J Physiol Endocrinol Metab;296(3):E445-453. 2009-01-01 [PMID: 19106247]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-57086

NB820-59212	Human Duodenum Whole Tissue Lysate (Adult Whole Normal)
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-57086

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



