Product Datasheet

CD79A Antibody (HM57) [DyLight 550] NB100-64347R

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-64347R

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB100-64347R



NB100-64347R

CD79A Antibody (HM57) [DyLight 550]

Product Information Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone HM57 Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate DyLight 550 Purity Protein G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 973 Gene Symbol CD79A Species Human Reactivity Notes Cross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information. Specificity/Sensitivity NB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expresed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b bole/blocking teps motion may intra signality following antigen-binding to sufface immunogloubulin. Clone HM57 has been reported to work in western blotting applications.		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneHM57Preservative0.05% Sodium AzideIsotypeIgG1ConjugateDyLight 550PurityProtein G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID973Gene SymbolCD79ASpecific Ty NotesHumanReactivity NotesCross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.Specificity/SensitivityNB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45KD protein that is expressed by B wmphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a nolecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor IBCR complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications.ImmunogenSynthetic peptide corresponding to 202-216 amino acid sequence of human mb-1NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone HM57 Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate DyLight 550 Purity Protein G purified Buffer 50mM Sodium Borate Product Description House Gene ID 973 Gene Symbol CD79A Species Human Reactivity Notes Cross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experimention. Specificity/Sensitivity NB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell traceptor ICRN complex. The CD79a AlcD79b heterodimers are also necessary for intracellular signalling following antigen-bioling to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications. Immunogen Synthetic peptide corresponding to 202-216 amino acid sequence of human mb-1 Notes	Unit Size	0.1 ml
ClonalMonoclonalCloneHM57Preservative0.05% Sodium AzideIsotypeIgG1ConjugateDyLight 550PurityProtein G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID973Gene SymbolCD79ASpeciesHumanReactivity NotesCross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse.Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.SpeciesNB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45KD protein that is sepressed by B lymphocyted during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications.ImmunogenSynthetic peptide corresponding to 202-216 amino acid sequence of human mb-1NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Concentration	•
CloneHM57Preservative0.05% Sodium AzideIsotypeIgG1ConjugateDyLight 550PurityProtein G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID973Gene SymbolCD79ASpeciesHumanReactivity NotesCross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.Specificity/SensitivityNB10-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin, Clone HM57 has been reported to work in western blotting applications.ImmunogenSynthetic peptide corresponding to 202-216 amino acid sequence of human mb-1NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate DyLight 550 Purity Protein G purified Buffer 50mM Sodium Borate Product Description Host Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Reactivity Notes Cross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information. Specificity/Sensitivity NB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b K29) to form a heterodimer that is non-covalently linked to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications. Immunogen Synthetic peptide corresponding to 202-216 amino acid sequence of human mb-1 Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Clonality	Monoclonal
Isotype IgG1 Conjugate DyLight 550 Purity Protein G purified Buffer 50mM Sodium Borate Product Description Host Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Reactivity Notes Cross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information. Specificity/Sensitivity NB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications. Immunogen Synthetic peptide corresponding to 202-216 amino acid sequence of human mb-1 Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Clone	HM57
Conjugate DyLight 550 Purity Protein G purified Buffer 50mM Sodium Borate Product Description Mouse Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Reactivity Notes Cross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information. Specificity/Sensitivity NB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signaling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications. Immunogen Synthetic peptide corresponding to 202-216 amino acid sequence of human mb-1 Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Preservative	0.05% Sodium Azide
Purity Protein G purified Buffer 50mM Sodium Borate Product Description Host Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Reactivity Notes Cross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information. Specificity/Sensitivity NB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications. Immunogen Synthetic peptide corresponding to 202-216 amino acid sequence of human mb-1 Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Isotype	IgG1
Buffer 50mM Sodium Borate Product Description Host Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Reactivity Notes Cross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information. Specificity/Sensitivity NB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications. Immunogen Synthetic peptide corresponding to 202-216 amino acid sequence of human mb-1 Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Conjugate	DyLight 550
Product Description Host Mouse Gene ID 973 Gene Symbol CD79A Species Human Reactivity Notes Cross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information. Specificity/Sensitivity NB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications. Immunogen Synthetic peptide corresponding to 202-216 amino acid sequence of human mb-1 Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Purity	Protein G purified
HostMouseGene ID973Gene SymbolCD79ASpeciesHumanReactivity NotesCross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.Specificity/SensitivityNB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications.ImmunogenSynthetic peptide corresponding to 202-216 amino acid sequence of human mb-1NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Buffer	50mM Sodium Borate
Gene ID 973 Gene Symbol CD79A Species Human Reactivity Notes Cross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information. Specificity/Sensitivity NB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications. Immunogen Synthetic peptide corresponding to 202-216 amino acid sequence of human mb-1 Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Product Description	
Gene Symbol CD79A Species Human Reactivity Notes Cross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information. Specificity/Sensitivity NB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications. Immunogen Synthetic peptide corresponding to 202-216 amino acid sequence of human mb-1 Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Host	Mouse
Species Human Reactivity Notes Cross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information. Specificity/Sensitivity NB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications. Immunogen Synthetic peptide corresponding to 202-216 amino acid sequence of human mb-1 Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Gene ID	973
Reactivity NotesCross reacts with Red Deer, Fallow Deer and American Bison. Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.Specificity/SensitivityNB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications.ImmunogenSynthetic peptide corresponding to 202-216 amino acid sequence of human mb-1NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Gene Symbol	CD79A
this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.Specificity/SensitivityNB100-64347 recognizes an epitope within the cytoplasmic domain of CD79a. CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications.ImmunogenSynthetic peptide corresponding to 202-216 amino acid sequence of human mb-1NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Species	Human
CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin. Clone HM57 has been reported to work in western blotting applications.ImmunogenSynthetic peptide corresponding to 202-216 amino acid sequence of human mb- 1NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Reactivity Notes	this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC
Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Specificity/Sensitivity	CD79a, also known as mb-1, is a 45kD protein that is expressed by B lymphocytes during differentiation from early pre-B cell stage through to plasma cells. The CD79a molecule associates with CD79b (B29) to form a heterodimer that is non-covalently linked to surface immunoglobulin, forming the B-cell receptor (BCR) complex. The CD79a/CD79b heterodimers are also necessary for intracellular signalling following antigen-binding to surface immunoglobulin.
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Immunogen	Synthetic peptide corresponding to 202-216 amino acid sequence of human mb- 1
Product Application Details	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
Applications Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin	Applications	Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-
Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, Immunohistochemistry- Frozen	Recommended Dilutions	
	Application Notes	Optimal dilution of this antibody should be experimentally determined.
	Application Notes	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NB100-64347R

NBP1-97005R	Mouse IgG1 Isotype Control (MG1) [DyLight 550]
NBP2-60209-50ug	Recombinant Human CD79A His Protein
7268-CT-100	CTLA-4 [Unconjugated]
9685-CD-050	CD79A [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-64347R

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

