Product Datasheet

COX-2 Antibody NB100-689

Unit Size: 0.5 ml

Store at 4C. Do not freeze.



Reviews: 1 Publications: 36

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-689

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB100-689



NB100-689


COX-2 Antibody

Product InformationUnit Size0.5 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C. Do not freeze.ClonalityPolyclonalPreservative0.05% Sodium AzideIsotypeIgGPurityAffinity purifiedBufferPBS (pH 7.4), 0.2% BSA, Tween-20Product DescriptionFreservativeHostRabbitGene SymbolST43SpeciesHuman, RatSpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenWestern Blot, Simple Western, Immunohistochemistry-Information).Product Application DetailsVestern Blot, Simple Western 1:100, Immunohistochemistry-Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot, Simple Western 1:100, Immunohistochemistry-Frozen reportedApplication NotesHC-P: recommended pretreatment of cirtate buffer, pH 6.0. Recommended in scientific literature (PMID: 31592103).		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C. Do not freeze.ClonalityPolyclonalPreservative0.05% Sodium AzideIsotypeIgGPurityAffinity purifiedBufferPBS (pH 7.4), 0.2% BSA, Tween-20Product DescriptionRabbitHostRabbitGene ID5743Gene SymbolPTGS2SpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsVestern Blot, Simple Western, Immunohistochemistry-Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry-FrozenApplication NotesHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Product Information	
services.StorageStore at 4C. Do not freeze.ClonalityPolyclonalPreservative0.05% Sodium AzideIsotypeIgGPurityAffinity purifiedBufferPBS (pH 7.4), 0.2% BSA, Tween-20Product DescriptionRabbitHostRabbitGene ID5743Gene SymbolPTGS2Specificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact arrino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry-Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry-Frozen reported incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Unit Size	0.5 ml
ClonalityPolyclonalPreservative0.05% Sodium AzideIsotypeIgGPurityAffinity purifiedBufferPBS (pH 7.4), 0.2% BSA, Tween-20Product DescriptionRabbitHostRabbitGene ID5743Gene SymbolPTGS2SpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry-frozen, Information 0:1:100, Immunohistochemistry-FrozenApplication NotesHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Concentration	
Preservative0.05% Sodium AzideIsotypeIgGPurityAffinity purifiedBufferPBS (pH 7.4), 0.2% BSA, Tween-20Product DescriptionPoduct DescriptionHostRabbitGene ID5743Gene SymbolPTGS2SpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry-Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry-FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Storage	Store at 4C. Do not freeze.
IsotypeIgGPurityAffinity purifiedBufferPBS (pH 7.4), 0.2% BSA, Tween-20Product DescriptionEtail (Construction)HostRabbitGene ID5743Gene SymbolPTGS2SpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry 1:50-1:100, Immunohistochemistry-FrozenApplication NotesHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Clonality	Polyclonal
PurityAffinity purifiedBufferPBS (pH 7.4), 0.2% BSA, Tween-20Product DescriptionHostRabbitGene ID5743Gene SymbolPTGS2SpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry-Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry-FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Preservative	0.05% Sodium Azide
BufferPBS (pH 7.4), 0.2% BSA, Tween-20Product DescriptionHostRabbitGene ID5743Gene SymbolPTGS2SpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry-FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Isotype	IgG
Product DescriptionHostRabbitGene ID5743Gene SymbolPTGS2SpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry 1:50-1:100, Immunohistochemistry-FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Purity	Affinity purified
HostRabbitGene ID5743Gene SymbolPTGS2SpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry-FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Buffer	PBS (pH 7.4), 0.2% BSA, Tween-20
Gene ID5743Gene SymbolPTGS2SpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry- Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry 1:50- 1:100, Immunohistochemistry-Paraffin 1:50-1:100, Immunohistochemistry- FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Product Description	
Gene SymbolPTGS2SpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry- Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry- FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Host	Rabbit
SpeciesHuman, RatSpecificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry- Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry 1:50- 1:100, Immunohistochemistry-Paraffin 1:50-1:100, Immunohistochemistry- FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Gene ID	5743
Specificity/SensitivityThis reacts with the 70 kDa protein.ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry- Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry 1:50- 1:100, Immunohistochemistry-Paraffin 1:50-1:100, Immunohistochemistry- FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Gene Symbol	PTGS2
ImmunogenRecombinant protein corresponding to the C-terminus of rat COX2 (the exact amino acid sequence is considered to be proprietary information).Product Application DetailsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry- Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry 1:50- 1:100, Immunohistochemistry-Paraffin 1:50-1:100, Immunohistochemistry- FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Species	Human, Rat
amino acid sequence is considered to be proprietary information).Product Application DetailsApplicationsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry- Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry 1:50- 1:100, Immunohistochemistry-Paraffin 1:50-1:100, Immunohistochemistry- FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Specificity/Sensitivity	This reacts with the 70 kDa protein.
ApplicationsWestern Blot, Simple Western, Immunohistochemistry, Immunohistochemistry- Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry 1:50- 1:100, Immunohistochemistry-Paraffin 1:50-1:100, Immunohistochemistry- FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Immunogen	
Frozen, Immunohistochemistry-ParaffinRecommended DilutionsWestern Blot 1:100-1:2000, Simple Western 1:100, Immunohistochemistry 1:50- 1:100, Immunohistochemistry-Paraffin 1:50-1:100, Immunohistochemistry- FrozenApplication NotesIHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Product Application Details	
1:100, Immunohistochemistry-Paraffin 1:50-1:100, Immunohistochemistry-Frozen Application Notes IHC-P: recommended pretreatment of citrate buffer, pH 6.0. Recommended incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Applications	
incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported	Recommended Dilutions	1:100, Immunohistochemistry-Paraffin 1:50-1:100, Immunohistochemistry-
	Application Notes	incubation time of 30 min at RT. Use in Immunohistochemistry-Frozen reported




Images

Immunohistochemistry-Paraffin: COX-2 Antibody [NB100-689] - Formalin fixed paraffin embedded colon carcinoma stained with COX-2 antibody (NB100-689).



Simple Western: COX-2 Antibody [NB100-689] - Simple Western lane view shows a specific band for COX2 in 1.0 mg/ml of Rat Brain at an antibody concentration of 1:100. This experiment was performed under reducing conditions using the 12-230 kDa separation system. Image provided through customer review.





Publications

Montecillo-Aguado M, Tirado-Rodriguez B, Antonio-Andres G et al. Omega-6 Polyunsaturated Fatty Acids Enhance Tumor Aggressiveness in Experimental Lung Cancer Model: Important Role of Oxylipins International Journal of Molecular Sciences 2022-05-31 [PMID: 35682855] (Immunohistochemistry)

Dib S, Loiola RA, Sevin E et al. TNF? Activates the Liver X Receptor Signaling Pathway and Promotes Cholesterol Efflux from Human Brain Pericytes Independently of ABCA1 International journal of molecular sciences 2023-03-22 [PMID: 36983062] (Western Blot, Human)

Lin FL, Cheng YW, Chen LH et al. Retinal protection by fungal product theissenolactone B in a sodium iodate-induced AMD model through targeting retinal pigment epithelial matrix metalloproteinase-9 and microglia activity Biomedicine & pharmacotherapie 2022-12-17 [PMID: 36535199] (WB, Mouse)

PeNa-Mercado E, Garcia-Lorenzana M, Huerta-Yepez S et al. Effect of melatonin on electrical impedance and biomarkers of damage in a gastric ischemia/reperfusion model PloS one 2022-08-16 [PMID: 35972989] (IHC-P, Rat)

Details:

Dilution used for IHC-P 1:1000

Jayakumar T, Yang CM, Yen TL et al. Anti-Inflammatory Mechanism of An Alkaloid Rutaecarpine in LTA-Stimulated RAW 264.7 Cells: Pivotal Role on NF-kappaB and ERK/p38 Signaling Molecules International journal of molecular sciences 2022-05-24 [PMID: 35682568] (WB, Mouse)

Ham HJ, Lee YS, Lee HP et al. G721-0282 Exerts Anxiolytic-Like Effects on Chronic Unpredictable Mild Stress in Mice Through Inhibition of Chitinase-3-Like 1-Mediated Neuroinflammation Frontiers in cellular neuroscience 2022-03 -07 [PMID: 35345530] (WB, Mouse)

Balde A, Raghavender P, Dasireddy S et al. Crab Pentapeptide and Its Anti-inflammatory Activity on Macrophage Cells International Journal of Peptide Research and Therapeutics 2021-09-08 (WB)

Eduardo P, Mario G, Carlos CEsar P et al. Bioelectric, tissue, and molecular characteristics of the gastric mucosa at different times of ischemia Experimental Biology and Medicine 2021-06-15 [PMID: 34130514]

Ham HJ, Lee YS, Yun J et al. K284-6111 alleviates memory impairment and neuroinflammation in Tg2576 mice by inhibition of Chitinase-3-like 1 regulating ERK-dependent PTX3 pathway Aging (Albany NY) 2020-11-07 [PMID: 33222690] (IF/IHC, Human)

Joshi I, Mohideen HS, Nazeer RA A Meretrix meretrix visceral mass derived peptide inhibits lipopolysaccharidestimulated responses in RAW264.7 cells and adult zebrafish model Int Immunopharmacol 2020-11-10 [PMID: 33187909]

Sukumaran V, Tsuchimochi H et al. Liraglutide Improves Renal Endothelial Function in Obese Zucker Rats on a High-Salt Diet. J Pharmacol Exp Ther 2019-01-06 [PMID: 30910920] (WB, Rat)

Jain S, Samal AG, Das B et al. Escherichia coli, a common constituent of benign prostate hyperplasia-associated microbiota induces inflammation and DNA damage in prostate epithelial cells Prostate 2020-08-24 [PMID: 32835423]

More publications at http://www.novusbio.com/NB100-689

www.novusbio.com





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NB100-689

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-689

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

