Product Datasheet

Acetyl-Lysine Antibody (1C6) NB100-74339

Unit Size: 100 ug

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 2 Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB100-74339

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB100-74339



NB100-74339

Acetyl-Lysine Antibody (1C6)


Product Information	
Unit Size	100 ug
Concentration	1 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	1C6
Preservative	0.05% Sodium Azide
Isotype	IgG1
Purity	Protein G purified
Buffer	PBS, 1 mg/mL BSA

Product Description	
Host	Mouse
Species	Non-species specific
Immunogen	Synthetic peptide sequence surrounding the acetylated lysine 9 of histone H3.
Product Application Details	


Product Application Details	
Applications	Western Blot, Dot Blot, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation, Chromatin Immunoprecipitation (ChIP)
	Western Blot 1:500 - 1:2000, Immunocytochemistry/ Immunofluorescence 1:100 - 1:500, Immunoprecipitation 3 ug, Dot Blot 1:100 - 1:2000, Chromatin Immunoprecipitation (ChIP) 10 ug

Images


Western Blot: Acetyl-Lysine Antibody (1C6) [NB100-74339] - Analysis of Acetyl-Lysine in HeLA cells.




Immunocytochemistry/Immunofluorescence: Acetyl-Lysine Antibody (1C6) [NB100-74339] - Analysis of lysine acetylated proteins (green) in HeLa cells left untreated (left panels) or treated with 0.3uM Trichostatin A for 16 hours (right panels).




Chromatin Immunoprecipitation: Acetyl-Lysine Antibody (1C6) [NB100-74339] - Analysis of chromatin proteins with Acetyl-Lysine was performed using cross-linked chromatin from 4-10^6 LNCaP cells. Immunoprecipitation was performed using the Magnetic ChIP kit with 10ug of an Acetyl-Lysine monoclonal antibody. Quantitation of immunoprecipitated GAPDH promoter sequence is presented as fold enrichment of the acetyl lysine monoclonal antibody versus non-specific IgG.



Western Blot: Acetyl-Lysine Antibody (1C6) [NB100-74339] - Analysis of Acetyl-Lysine was performed by loading various amounts of acetylated BSA or non-acetylated BSA per well and 10ul of PageRuler Prestained Protein Ladder onto a 4-20% Tris-HCl polyacrylamide gel. Proteins were transferred to a PVDF membrane and blocked with 5% BSA/TBST for at least 1 hour. The membrane was probed with an Acetyl-Lysine monoclonal antibody at a dilution of 1:1000 overnight at 4C on a rocking platform, washed in TBS-0.1%Tween-20, and probed with a goat antimouse IgG-HRP secondary antibody at a dilution of 1:20,000 for 1 hour. Chemiluminescent detection was performed using SuperSignal West Pico.




Western Blot: Acetyl-Lysine Antibody (1C6) [NB100-74339] - Analysis of lysine acetylated proteins from cells left untreated (DMSO only) or cells treated with 0.3uM or 3uM of Trichostatin A (TSA) for 16 hours was performed by loading 50 ug of the indicated whole cell lysates per well and 10ul of PageRuler Prestained Protein Ladder onto a 4-20% Tris-HCl polyacrylamide gel. Proteins were transferred to a PVDF membrane and blocked with 5% BSA/TBST for at least 1 hour. The membrane was probed with an Acetyl-Lysine monoclonal antibody at a dilution of 1:1000 overnight at 4C on a rocking platform, washed in TBS-0.1%Tween-20, and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:20,000 for 1 hour. Chemiluminescent detection was performed using SuperSignal West Pico.




Dot Blot: Acetyl-Lysine Antibody (1C6) [NB100-74339] - Specific detection of lysine acetylated histones using an Acetyl-Lysine monoclonal antibody. Histone peptides containing 384 different modification combinations spotted in duplicate on a peptide array were used according to the manufacturer instructions. A schematic and table with representative results (positive detection in red, negative in blue) are shown. Result interpretation and quantification were performed using the manufacturer software. Chemiluminescent detection was performed.

Ш	A A P	position	Histone peptide sequence (N->C)	result
Ш	× 2 0	AL.	DHARTKOTARESTOGEAPRED	- 0
Ш	76.6 H3 57.6 H3 H3	A2	HOLARING TAGTAREST GGEAPREG	- 4
Ш	R KRAK S T G G Historie H3] R KRAK S T G G Q [Historie H3] Q T A R K S T G	Wil	(HS) A EXCHIRGT A REST G G KAPREG	15
Ш	2TA R KBas S EKQ (Histona 2TA R KBas S R KQ (Histon R KQ (Histon R K Q (Histon	AS:	HUHAR THE KOT ARKST GGEAPHED	194
Ш	40 42 50	All	DEDARTERNOQUARESTGGEARREQ	-
Ш	7 2	A9	MRAKIHALQTAKKSTODEAPHKQ	122
	7 5 5	Atti	OHDARIEQ SARKWEST GGRAFREQ	
	1 1 1	CZ.	IHEDARTKQTARIRWEIKSSESTGGKAPHKQ	2.0
1	1=1=1=	7 (8	(H3)ARTKGTARRoots KlocsTGGKAPHKG	+
	, ,	014	(H3)ARTEGTACIEMESTGGEAPEED	
		04	[HXDARTKOTAR KRAIST G G KLABOCA PREC	***
	88888888888888888888888888	8 £24	(HS) A House I Hamel Q TAR KNess TO GRAPHEQ	
		84	OHIGA RETRAINED QUANTITIES STEER APPRICA	
	STOCK CHARGE CONTROL	611	HERART MADE OF ANTHROLD MESONS TO GRAPH K.Q.	**
Array rows	0.000	а.	POIFEKGLATKAAFEZZALSAFATGG	***
Tay.	(C.	118	DESIPREGLATE A A Rismon 827ac SA PATG G	***
A	(I)	8 MII	HEIS GROKSOCO GREEN GLORIZANG GREATA	+++
1	003707000700000000000000000000000000000	N4	HKIGKGGAKNHRN20scVLRDN/QG/T	***
400	Array columns 1-24	P16	(HUB) PO PARSec SAPA PER LINE GITTER RISEC PAY T	444


Immunoprecipitation: Acetyl-Lysine Antibody (1C6) [NB100-74339] - Immunoprecipitation of acetylated alpha-Tubulin was performed.3uM or 3uM Trichostatin A (TSA) for 16 hours. Antigen-antibody complexes were formed by incubating 500ug of lysate with 3ug of an Acetyl-Lysine monoclonal antibody overnight on a rocking platform at 4C. The immune complexes were captured on 50ul Protein A/G Agarose at a dilution of 1:1000 overnight rotating at 4C, washed in TBST, and probed with Clean-blot IP Detection Reagent at a dilution of 1:2000 for at least 1 hour.



Immunoprecipitation: Acetyl-Lysine Antibody (1C6) [NB100-74339] - Analysis of acetylated Histone H3 (Lys9) was performed using whole cell lysates from cells left untreated (DMSO only) or cells treated with 0.3uM or 3uM Trichostatin A (TSA) for 16 hours. Antigen-antibody complexes were formed by incubating 500ug of the indicated lysate with 3ug of an Acetyl-Lysine mAb overnight on a rocking platform at 4C. The immune complexes were captured on 50ul Protein A/G Agarose, washed extensively, and eluted with 5X Lane Marker Reducing Sample Buffer. Samples were resolved on a 4-20% Tris-HCl polyacrylamide gel, transferred to a PVDF membrane, and blocked with 5% BSA/TBS-0.1%Tween for at least 1 hr. The membrane was probed with an Acetyl-Histone H3 (Lys9) mAb at a dilution of 1:1000 overnight rotating at 4C, washed in TBST, and probed with Clean-blot IP Detection Reagent at a dilution of 1:2000 for at least 1 hour. Chemiluminescent detection was performed using SuperSignal West Pico.



Dot Blot: Acetyl-Lysine Antibody (1C6) [NB100-74339] - Specific detection of lysine acetylated histones using an Acetyl-Lysine monoclonal antibody. Histone peptides containing 384 different modification combinations spotted in duplicate on a peptide array were used according to the manufacturer's instructions (Active Motif). Representative results showing specific detection of acetylated Histone H4 K20 (blue circles, panel A) and H3 K9 (blue circles, panel B). Note that Histone H4 K20me2/K20me3 (red and yellow circles, panel A) and H3 K9me2/K9m3 (red and yellow circles, panel B) are not detected by the Acetyl-Lysine monoclonal antibody.



Publications

Dobosz AM, Janikiewicz J, Borkowska AM et al. Stearoyl-CoA Desaturase 1 Activity Determines the Maintenance of DNMT1-Mediated DNA Methylation Patterns in Pancreatic ?-Cells International Journal of Molecular Sciences 2020-09-18 [PMID: 32961871]

Dziewulska A, Dobosz AM, Dobrzyn A et al. SCD1 regulates the AMPK/SIRT1 pathway and histone acetylation through changes in adenine nucleotide metabolism in skeletal muscle J. Cell. Physiol. 2019-06-26 [PMID: 31241768] (Mouse)

Tataranni T, Agriesti F, Ruggieri V et al. Rewiring carbohydrate catabolism differentially affects survival of pancreatic cancer cell lines with diverse metabolic profiles Oncotarget 2017-06-20 [PMID: 28476035] (WB, Human)





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB100-74339

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-97005-0.5mg Mouse IgG1 Isotype Control (MG1)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-74339

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

