

Product Datasheet

N myc interactor Antibody - BSA Free NB100-763

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-763

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-763



NB100-763

N myc interactor Antibody - BSA Free

Product Information	
Unit Size	100 ul
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)


Product Description	
Host	Rabbit
Gene ID	9111
Gene Symbol	NMI
Species	Human
Immunogen	The immunogen recognized by this antibody maps to a region between residues 1 and 50 of human N-myc and STAT interactor using the numbering given in entry NP_004679.1 (GeneID 9111).

Product Application Details	
Applications	Immunohistochemistry, Immunoprecipitation, Western Blot (Negative)
Recommended Dilutions	Immunohistochemistry, Immunoprecipitation 2-5 ug/mg lysate, Western Blot (Negative)
Application Notes	Immunoprecipitation Not Recommended for Western Blot. Use rabbit anti-NMI antibody, Cat. No. NB 100-764.




Images

Immunohistochemistry: N myc interactor Antibody [NB100-763] -
 Sample: FFPE section of human ovarian carcinoma. Antibody: Affinity
 purified rabbit anti- NMI used at a dilution of 1:1,000 (1ug/ml). Detection:
 DAB



Immunoprecipitation: N myc interactor Antibody [NB100-763] - Detection
 of human NMI/N myc interactor by western blot of immunoprecipitates.
 Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP
 loaded) from HeLa cells prepared using NETN lysis buffer. Antibodies:
 Affinity purified rabbit anti-NMI antibody NB100-763 (lot 2) used for IP at
 6 ug per reaction. NMI was also immunoprecipitated by rabbit anti-NMI
 antibody NB100-763-1. For blotting immunoprecipitated NMI, NB100-763
 was used at 1 ug/ml. Detection: Chemiluminescence with an exposure
 time of 30 seconds.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-763

NBL1-13690	N myc interactor Overexpression Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-763

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

