

Product Datasheet

MELK Antibody NB100-782

Unit Size: 0.1 mg

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-782

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-782



NB100-782

MELK Antibody


Product Information	
Unit Size	0.1 mg
Concentration	0.5 mg/ml
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris saline (20 mM Tris pH 7.3, 150 mM NaCl), 0.5% BSA

Product Description	
Host	Goat
Gene ID	9833
Gene Symbol	MELK
Species	Human
Immunogen	Peptide with sequence KRLVEDILSSCKV corresponding to C-Terminus according to NP_055606.1.


Product Application Details	
Applications	Western Blot, Peptide ELISA
Recommended Dilutions	Western Blot 0.1 - 0.3 ug/ml, Peptide ELISA Detection limit 1:32000
Application Notes	WB: Experiments gave bands at approx. 75 kDa band and 150 kDa band in human testis lysates and in cell lines HeLa and Jurkat after 0.1 ug/ml antibody staining. The 150 kDa band is explained by glycosylation of this member of the AMPK family. This protein has a calculated MW of 74.6 kDa band according to NP_055606.1.

Images


Western Blot: MELK Antibody [NB100-782] - MELK in human testes lysate (RIPA buffer, 1.4E5 cells per lane) using this antibody at 0.1 ug/ml.



Western Blot: MELK Antibody [NB100-782] - NB100-782 (0.1ug/ml) staining of HEPG2 cell lysate (35ug protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence. Online Catalogue NAC1 to Quaking > Goat Anti-PPP1R15A Antibody



Western Blot: MELK Antibody [NB100-782] - NB 100-782 (0.5ug/ml) staining of Jurkat lysate (RIPA buffer, 1.4E5 cells per lane). Detected by western blot using chemiluminescence.



Publications

Han AR, Lee JY, Kim HJ et al. A CXCR4 antagonist leads to tumor suppression by activation of immune cells in a leukemia-induced microenvironment. *Oncol. Rep.* 2015-09-21 [PMID: 26398122]

Nagase T, Seki N, Ishikawa K et al. Prediction of the coding sequences of unidentified human genes. V. The coding sequences of 40 new genes (KIAA0161-KIAA0200) deduced by analysis of cDNA clones from human cell line KG-1. *DNA Res* 1996 Feb 29;3(1):17-24 1996-02-01 [PMID: 8724849]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-782

NB800-PC1	HeLa Whole Cell Lysate
HAF017	Rabbit anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
HAF109	Donkey anti-Goat IgG Secondary Antibody [HRP (Horseradish Peroxidase)]
NB410-28088-1mg	Goat IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-782

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications



