

Product Datasheet

PHF8 Antibody NB100-93314

Unit Size: 0.1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB100-93314

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB100-93314



NB100-93314

PHF8 Antibody

Product Information	
Unit Size	0.1 ml
Concentration	0.2 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	TBS and 0.1% BSA


Product Description	
Host	Rabbit
Gene ID	23133
Gene Symbol	PHF8
Species	Human
Immunogen	The immunogen recognized by this antibody maps to a region between residue 925 and 975 of human PHD finger protein 8 using the numbering given in entry CAI41576.1 (GeneID 23133).

Product Application Details	
Applications	Western Blot, Immunoprecipitation, Knockdown Validated
Recommended Dilutions	Western Blot 1:2000-1:10000, Immunoprecipitation, Knockdown Validated
Application Notes	Use in immunoprecipitation reported in scientific literature (PMID: 27991916). Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.




Images


Western Blot: PHF8 Antibody [NB100-93314] - Detection of Human PHF8 by Western Blot. Samples: Whole cell lysate (50 ug) from HeLa, HeLa (RIPA), 293T, and Jurkat cells. Antibodies: Affinity purified rabbit anti-PHF8 antibody NB100-93314 used for WB at 0.04 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.



Knockdown Validated: PHF8 Antibody [NB100-93314] - Western blot analysis showing knockdown of PHF8 by shRNA against PHF8. LNCaP cells were infected with a lentiviral shPHF8 and western blot analysis was carried out with anti-PHF8 antibody. Image collected and cropped by CiteAb from the following publication ([nature.com/articles/oncsis201674](https://www.nature.com/articles/oncsis201674)), licensed under a CC-BY license.



Western Blot: PHF8 Antibody [NB100-93314] - PHF8 interacts with the AR & functions as an AR coactivator. (a) PHF8 interacts with AR independent of DHT. HEK293T cells were transiently transfected with plasmids expressing Flag-tagged PHF8 & GFP-tagged AR. Cells were treated with or without 10 nM DHT for 48 h & the cell lysates were used for immunoprecipitation with antibodies against either Flag or GFP. Western blotting was conducted with antibodies as indicated. (b) PHF8 interacts with AR independent of its demethylase activity. HEK293T cells were transiently transfected with plasmids expressing Flag-tagged PHF8 (H247A) mutant & GFP-tagged AR. The interaction assay by co-immunoprecipitation was performed as described above. (c) In LNCaP cells endogenous PHF8 interacts with AR. LNCaP cells were treated with 10 nM DHT for 48 h & the cell lysates were used for immunoprecipitation with antibodies against either PHF8 or AR. Western blotting was conducted with AR or PHF8 antibody as indicated. (d) PHF8 enhances AR transcriptional activation in a demethylase activity-dependent manner. HeLa cells were co-transfected with the 4 × UAS-TK-luc reporter, the plasmids expressing the fusion protein Gal-AR, & increased amount of either the wild-type PHF8 or mutant PHF8 (H247A) as indicated. Cells were cultured with or without 10 nM R1881 overnight. The relative luciferase activities in the cell lysates were normalized to renilla luciferase activity. The relative luciferase activities are presented as mean±s.e.m. of three independent transfections & the levels of PHF8 were estimated by western blot as depicted in the bottom panel. Image collected & cropped by CiteAb from the following publication (<https://www.nature.com/articles/oncsis201674>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Tong D, Liu Q, Liu G et al. The HIF/PHF8/AR axis promotes prostate cancer progression. *Oncogenesis*. 2016-12-19 [PMID: 27991916] (IP, Human)

Zhu Z, Wang Y, Li X et al. PHF8 is a histone H3K9me2 demethylase regulating rRNA synthesis. *Cell Res* 2010-07-01 [PMID: 20531378]





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB100-93314

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP2-56147PEP	PHF8 Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB100-93314

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

