

Product Datasheet

Abhd5 Antibody - BSA Free

NB110-41576

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 29

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB110-41576

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB110-41576



NB110-41576

Abhd5 Antibody - BSA Free

Product Information

Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS

Product Description

Description	Novus Biologicals Rabbit Abhd5 Antibody - BSA Free (NB110-41576) is a polyclonal antibody validated for use in WB and Simple Western. Anti-Abhd5 Antibody: Cited in 29 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	51099
Gene Symbol	ABHD5
Species	Human, Mouse, Rat
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID:23878361)
Immunogen	A synthetic peptide made to an internal region within residues 200-300 of the human Abhd5 protein. [Swiss-Prot# Q8WTS1]


Product Application Details

Applications	Western Blot, Simple Western
Recommended Dilutions	Western Blot 1:500, Simple Western 1:1000
Application Notes	<p>A band is seen at ~43 kDa in Western Blot. In ICC/IF cytoplasmic staining was observed in HeLa cells.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.</p> <p>See Simple Western Antibody Database for Simple Western validation: Tested in MCF-7 lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:1000, apparent MW was 55 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.</p>




Images


Western Blot: Abhd5 Antibody [NB110-41576] - CGI-58 western blot protein bands compared to representative Ponceau bands as equal loading control. Image collected and cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0120136>), licensed under a CC-BY license.




Western Blot: Abhd5 Antibody [NB110-41576] - Abhd5 overexpression lysates (NBL1-07199).



Western Blot: Abhd5 Antibody [NB110-41576] - Factors regulating lipid storage and breakdown in C2C12 muscle cells. Muscle cells were incubated with PALM (open), NORM (filled), or HSFA (hatched). Protein abundance of ATGL. In all figure panels data are expressed relative to a no fatty acid control condition (0mM). In panel B, #P<0.05 for a main effect of HSFA vs. PALM and NORM. Representative blots are inset above each figure panel. GPAT, glycerol-3-phosphate acyltransferase; DGAT, diacylglycerol acyltransferase; ATGL, adipose triacylglycerol lipase; CGI-58, comparative gene identification 58; HSL, hormone sensitive lipase; AU, arbitrary units. Image collected and cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0120871>), licensed under a CC-BY license.



Western Blot: Abhd5 Antibody [NB110-41576] - MCF7 cell lysates.




Simple Western: Abhd5 Antibody [NB110-41576] - Image shows a specific band for Abhd5 in 0.5 mg/mL of MCF-7 lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.




Protein content of important players involved in lipid droplet dynamics in Caucasian and South Asian subjects before and after a 5-day HFHC-diet. (a) ATGL, (b) CGI-58, (c) PLIN2, (d) PLIN3 and (e) PLIN5 protein content. Data are presented as mean \pm SEM and were statistically analyzed with a Repeated Measures ANOVA; * $P < 0.05$ for diet effect, # $P < 0.05$ for group effect.

b




The turnover of DAG and TAG is increased in the DGAT1 overexpressing TA muscle. (A) Western blotting of ATGL, (B) CGI58 (C) and ADRP in rat TA muscle. Data are expressed as mean \pm SEM ($n = 10-12$). # $P < 0.05$ HFD-DGAT1 vs. HFD-control.

B




Western Blot: Abhd5 Antibody - BSA Free [NB110-41576] - A. ATGL ($n = 8$) & CGI-58 ($n = 7$) protein content in the three skeletal muscles (SOL, RG, & WG & heart). Ponceau stain for total protein was used as a loading control. Data are reported as mean \pm SE & bars with the same letter are not significantly different within the skeletal muscles ($p < 0.05$). Insets: representative blots: lane 1, WG (white gastrocnemius); lane 2, RG (red gastrocnemius); lane 3, SOL (soleus). B. ATGL & CGI-58 western blot protein bands compared to representative Ponceau bands as equal loading control. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0120136>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.


B.




Western Blot: Abhd5 Antibody - BSA Free [NB110-41576] - The turnover of DAG & TAG is increased in the DGAT1 overexpressing TA muscle. (A) Western blotting of ATGL, (B) CGI58 (C) & ADRP in rat TA muscle. Data are expressed as mean \pm SEM (n=10–12). # $P < 0.05$ HFD-DGAT1 vs. HFD-control. Image collected & cropped by CiteAb from the following publication (<https://dx.plos.org/10.1371/journal.pone.0014503>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: Abhd5 Antibody - BSA Free [NB110-41576] - Effects of treadmill running on the mRNA & protein expressions of ATGL (A,E), CGI-58 (B,F), G0S2 (C,G), HSL (D,H) in the right ventricle. Data are expressed as mean \pm SEM. For the sake of clarity, the control group was set at 100, & exercised groups were scaled with respect to Ctrl * $p < 0.05$ difference vs. control (Ctrl); # $p < 0.05$ difference M120 vs. M30; \$ $p < 0.05$ difference F30 vs. M30. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31137663>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: Abhd5 Antibody - BSA Free [NB110-41576] - Protein content of important players involved in lipid droplet dynamics in Caucasian & South Asian subjects before & after a 5-day HFHC-diet. (a) ATGL, (b) CGI-58, (c) PLIN2, (d) PLIN3 & (e) PLIN5 protein content. Data are presented as mean \pm SEM & were statistically analyzed with a Repeated Measures ANOVA; * $P < 0.05$ for diet effect, # $P < 0.05$ for group effect. Image collected & cropped by CiteAb from the following publication (<https://www.nature.com/articles/srep42393>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: Abhd5 Antibody - BSA Free [NB110-41576] - Effects of treadmill running on the mRNA & protein expressions of ATGL (A,E), CGI-58 (B,F), G0S2 (C,G), HSL (D,H) in the left ventricle. Data are expressed as mean \pm SEM. For the sake of clarity, the control group was set at 100, & exercised groups were scaled with respect to Ctrl * $p < 0.05$ difference vs. control (Ctrl); # $p < 0.05$ difference M120 vs. M30; \$ $p < 0.05$ difference F30 vs. M30. Adipose triglyceride lipase (ATGL), comparative gene identification-58 (CGI-58), G0/G1 switch gene 2 (G0S2), hormone sensitive lipase (HSL). Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31137663>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Bao X, Ma X, Huang R et al. Knockdown of hepatocyte Perilipin-3 mitigates hepatic steatosis and steatohepatitis caused by hepatocyte CGI-58 deletion in mice *Journal of Molecular Cell Biology* 2022-12-26 [PMID: 36107452]

Sundararajan Mahalingam, Ramesh Bellamkonda, Kusum K Kharbanda, Madan Kumar Arumugam, Vikas Kumar, Carol A Casey, Lorenzo Leggio, Karuna Rasineni Role of ghrelin hormone in the development of alcohol-associated liver disease. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie* 2024-04-18 [PMID: 38640709]

Mahalingam S, Bellamkonda R, Arumugam MK et al. Glucagon-like peptide 1 receptor agonist, exendin-4, reduces alcohol-associated fatty liver disease *Biochemical pharmacology* 2023-05-18 [PMID: 37209859] (WB, Rat)

Chrzanowski-Smith OJ, Edinburgh RM, Smith E et al. Resting skeletal muscle PNPLA2 (ATGL) and CPT1B are associated with peak fat oxidation rates in men and women but do not explain observed sex differences *Experimental physiology* 2021-03-06 [PMID: 33675111] (WB, Human)

Fievet A, Bellanger D, Rieunier G et al. Assessment of the Main Compounds of the Lipolytic System in Treadmill Running Rats: Different Response Patterns between the Right and Left Ventricle *Int J Mol Sci* 2019-05-24 [PMID: 31137663] (WB, Rat)

Daemen S, Gemmink A, Brouwers B et al. Distinct lipid droplet characteristics and distribution unmask the apparent contradiction of the athlete's paradox *Molecular Metabolism* 2018-08-01 [PMID: 30174227] (WB, Human)

Rogne M, Chu DT, Kuntziger TM et al. OPA1-anchored PKA phosphorylates perilipin 1 on S522 and S497 in adipocytes differentiated from human adipose stem cells. *Mol. Biol. Cell* 2018-04-24 [PMID: 29688805] (Human)

Perreault L, Newsom SA, Strauss A, Kerege A. Intracellular localization of diacylglycerols and sphingolipids influences insulin sensitivity and mitochondrial function in human skeletal muscle. *JCI Insight*. 2018-02-08 [PMID: 29415895] (WB, Human)

Snook LA, Trottier SK, Worndl EA et al. Prior Endurance Training Enhances Beta-Adrenergic Signaling in Epididymal Adipose from Mice Fed a High-Fat Diet Obesity (Silver Spring) 2017-10-01 [PMID: 28857453] (Mouse)

Gemmink A, Bakker LE, Guigas B et al. Lipid droplet dynamics and insulin sensitivity upon a 5-day high-fat diet in Caucasians and South Asians. *Sci Rep*. 2017-02-14 [PMID: 28195217] (WB, Human)

Turnbull PC, Longo AB, Ramos SV et al. Increases in skeletal muscle ATGL and its inhibitor G0S2 following 8 weeks of endurance training in metabolically different rat skeletal muscles. *Am. J. Physiol. Regul. Integr. Comp. Physiol*. 2015-10-28 [PMID: 26511521] (WB, Rat)

Sitnick MT, Basantani MK, Cai L et al. Skeletal muscle triacylglycerol hydrolysis does not influence metabolic complications of obesity. *Diabetes* 2013-10-01 [PMID: 23835334] (Mouse)

More publications at <http://www.novusbio.com/NB110-41576>



Procedures

Western Blot protocol for Abhd5 Antibody (NB110-41576)

Western Blot Protocol

1. Perform SDS-PAGE (4-12% MOPS) on samples to be analyzed, loading 40 ug of total protein per lane.
2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
3. Rinse membrane with dH₂O and then stain the blot using Ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
4. Rinse the blot in TBS for approximately 5 minutes.
5. Block the membrane using 5% non-fat dry milk + 1% BSA in TBS, 1 hour at room temperature.
6. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
7. Dilute the rabbit anti-CGI58 primary antibody (NB 110-41576) in blocking buffer and incubate 2 hours at room temperature.
8. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
9. Apply the diluted rabbit-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturer's instructions) and incubate 1 hour at room temperature.
10. Wash the blot in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturer's instructions (Pierce ECL).

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.

Immunocytochemistry/Immunofluorescence protocol for Abhd5 Antibody (NB110-41576)

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.
4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,000 and incubate for 10 minutes. Wash a third time for 10 minutes.
9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB110-41576

NB820-59253	Human Skeletal Muscle Whole Tissue Lysate (Adult Whole Normal)
NB110-41576PEP	Abhd5 Antibody Blocking Peptide
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB110-41576

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

