

Product Datasheet

CD36 Antibody (D-2712) - BSA Free NB110-59724

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 **Publications: 12**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB110-59724

Updated 10/23/2024 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB110-59724




NB110-59724

CD36 Antibody (D-2712) - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	D-2712
Preservative	0.05% Sodium Azide
Isotype	IgA Kappa
Purity	Protein G purified
Buffer	PBS
Target Molecular Weight	110 kDa
Product Description	
Host	Mouse
Gene ID	948
Gene Symbol	CD36
Species	Mouse, Rat, Human (Negative)
Reactivity Notes	Does not react with human.
Marker	Endothelial Cell Marker
Immunogen	This CD36 Antibody (D-2712) was developed against adenovirus expressing recombinant mouse CD36 [Uniprot: Q08857]
Product Application Details	
Applications	Western Blot, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunoprecipitation
Recommended Dilutions	Western Blot 1:500, Immunohistochemistry 1:100-1:250, Immunoprecipitation 10-20 ug, Immunohistochemistry-Paraffin 1:100-1:250
Application Notes	In Western blot, a band is observed at ~70-80 kDa. The theoretical molecular weight of CD36 is ~53 kDa. The difference in theoretical MW and actual MW, as seen in Western blot, is most likely due to the heavy glycosylation and palmitoylation of this protein.

Images

Western Blot: CD36 Antibody (D-2712) [NB110-59724] - Detection of CD36 in mouse adipose lysate.



Immunohistochemistry: CD36 Antibody (D-2712) [NB110-59724] - Staining of CD36 in mouse liver.



Publications

Liang Y, Li X, Zhang Y et al. Induced Pluripotent Stem Cells-Derived Mesenchymal Stem Cells Attenuate Cigarette Smoke-Induced Cardiac Remodeling and Dysfunction *Front Pharmacol* 2017-08-14 [PMID: 28804458]

Ji Y, Liang Y, Chu PH Et al. The effects of intermittent hypoxia on hepatic expression of fatty acid translocase CD36 in lean and diet-induced obese mice *Biomed J* 2022-10-16 [PMID: 36244649] (WB, Mouse)

Details:

Citation using the HRP version of this antibody.

Georgiou, D K, Dagnino-Acosta, A Et al. Ca²⁺ Binding/Permeation via Calcium Channel, CaV1.1, Regulates the Intracellular Distribution of the Fatty Acid Transport Protein, CD36, and Fatty Acid Metabolism. *J Biol Chem* 2015-09-25 [PMID: 26245899] (IF/IHC, Mouse)

Details:

Citation using the DyLight 550 format of this antibody.

Yuan C, Guo X, Zhou Q et al. OAB-14, a bexarotene derivative, improves Alzheimer's disease-related pathologies and cognitive impairments by increasing beta-amyloid clearance in APP/PS1 mice. *Biochim Biophys Acta Mol Basis Dis* 2019-01-01 [PMID: 30389579] (Mouse)

Kuhnel E, Kleff V, Stojanovska V et al. Placental Specific Overexpression of sFlt1 Alters Trophoblast Differentiation and Nutrient Transporter Expression in an IUGR Mouse Model. *Journal of Cellular Biochemistry*. 2016-11-09 [PMID: 27859593] (WB, Mouse)

Nusken E, Gellhaus A, Kuhnel E et al. Increased Rat Placental Fatty Acid, But Decreased Amino Acid and Glucose Transporters Potentially Modify Intrauterine Programming. *J. Cell. Biochem*. 2015-11-21 [PMID: 26590355] (WB, Rat)

Chen YJ, Lo YH, Chen YT et al. Magnesium lithospermate B improves metabolic changes in high-fat diet-fed rats with metabolic syndrome *Journal of Functional Foods*. 2015-04-01 (WB, Rat)

Details:

CD36/SR-B3 antibody was used for WB application on lysates of liver tissue from rats that were fed different diet regimens as - normal diet control/NC, high-fat diet control/HC or HC+ MLB/Magnesium lithospermate B (MLB5-20/HC +5-20?mg/kg/day of MLB). Primary antibody was used at 1:1000 dilution in 5% skim milk and detected with HRP-conjugated secondary antibody - chemiluminescent HRP substrate (data shown in Fig. 4A).

Gong T, Wang Q, Lin Z et al. Endoplasmic reticulum (ER) stress inhibitor salubrinal protects against ceramide-induced SH-SY5Y cell death. *Biochem Biophys Res Commun*. 2012-01-01 [PMID: 22935424]

Feldhahn N, Arutyunyan A, Stoddart S et al. Environment-mediated drug resistance in Bcr/Abl-positive acute lymphoblastic leukemia *Oncoimmunology* 2012-08-01 [PMID: 22934254] (WB, Mouse)

During A, Doraiswamy S, Harrison EH. Xanthophylls are preferentially taken up compared with {beta}-carotene by retinal cells via a SRBI-dependent mechanism. *J Lipid Res*;49(8):1715-1724. 2008-01-01 [PMID: 18424859] (WB, Human)

Febbraio M et al. Targeted disruption of the class B scavenger receptor CD36 protects against atherosclerotic lesion development in mice. *J Clin Invest*;105(8):1049-56. 2000-04-01 [PMID: 10772649] (IF/IHC, Mouse)

Duan SZ, Ivashchenko CY, Russell MW, Milstone DS, Mortensen RM. Cardiomyocyte-specific knockout and agonist of peroxisome proliferator-activated receptor-gamma both induce cardiac hypertrophy in mice. *Circ Res*;97(4):372-9. 2005-08-19 [PMID: 16051889] (WB, IF/IHC, Mouse)

Procedures

Western Blot protocol for CD36 Antibody (NB110-59724)

CD36 Antibody (D-2712):

1. Perform SDS-PAGE (4-12% MOPS) on samples to be analyzed, loading 30 ug of total protein per lane.
 2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
 3. Rinse membrane with dH₂O and then stain the blot using Ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
 4. Rinse the blot in TBS for approximately 5 minutes.
 5. Block the membrane using 5% non-fat dry milk + 1% BSA in TBS, 1 hour at room temperature.
 6. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
 7. Dilute the rabbit anti-CD36 primary antibody (NB 110-59724) in blocking buffer and incubate 2 hours at room temperature.
 8. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
 9. Apply the diluted mouse-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturer's instructions) and incubate 1 hour at room temperature.
 10. Wash the blot in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each (this step can be repeated as required to reduce background).
 11. Apply the detection reagent of choice in accordance with the manufacturer's instructions (Pierce, ECL).
- **Note:** Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.

Immunohistochemistry-Paraffin protocol for CD36/SR-B3 Antibody (NB110-59724)

CD36 Antibody (D-2712):

Antigen Unmasking:

Bring slides to a boil in 10mM sodium citrate buffer (pH 6.0) then maintain at a sub-boiling temperature for 10 minutes. Cool slides on bench top for 30 minutes.

Staining:

1. Wash sections in dH₂O three times for 5 minutes each.
2. Wash section in wash buffer (1X PBS/0.1% Tween-20 (1X PBST)) for 5 minutes.
3. Block each section with 100-400 ul blocking solution (1X PBST, 5% goat serum) for 1 hour at room temperature.
4. Remove blocking solution and add 100-400 ul primary antibody diluted in 1X PBST, 5% goat serum to each section. Incubate overnight at 4C.
5. Remove antibody solution and wash sections in wash buffer three times for 5 minutes each.
6. Add 100-400 ul biotinylated secondary antibody, diluted in 1X PBST, 5% goat serum. Incubate 30 minutes at room temperature.
7. Remove secondary antibody solution and wash sections three times with wash buffer for 5 minutes each.
8. Add 100-400 ul Streptavidin-HRP reagent to each section and incubate for 30 minutes at room temperature.
9. Wash sections three times in wash buffer for 5 minutes each.
10. Add 100-400 ul DAB substrate to each section and monitor staining closely.
11. As soon as the sections develop, immerse slides in dH₂O.
12. Counterstain sections in hematoxylin.
13. Wash sections in dH₂O two times for 5 minutes each.
14. Dehydrate sections.
15. Mount the coverslips.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB110-59724

NBL1-08939	CD36 Overexpression Lysate
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NB110-59724H	CD36 Antibody (D-2712) [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB110-59724

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

