

Product Datasheet

VPS34 Antibody NB110-87320

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 8


Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB110-87320

Updated 2/17/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB110-87320



NB110-87320**VPS34 Antibody**


Product Information	
Unit Size	0.1 ml
Concentration	0.2 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS, 0.1% BSA, and 50% Glycerol
Target Molecular Weight	100 kDa
Product Description	
Host	Rabbit
Gene ID	5289
Gene Symbol	PIK3C3
Species	Human, Mouse, Rat, Porcine
Reactivity Notes	Immunogen displays the following percentage of sequence identity for non-tested species: Xenopus (87%).
Immunogen	Synthetic peptide made to an internal portion of the human VPS34 protein (within residues 700-850). [Swiss-Prot# Q8NEB9]
Product Application Details	
Applications	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Knockdown Validated
Recommended Dilutions	Western Blot 1:1000, Simple Western 1:20, Immunocytochemistry/ Immunofluorescence 1:100, Knockdown Validated
Application Notes	<p>This VPS34 antibody is useful for Immunocytochemistry/Immunofluorescence and Western blot, where a band is seen approx. 100 kDa.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.</p> <p>See Simple Western Antibody Database for Simple Western validation: Tested in HepG2 lysate 0.5 mg/mL, separated by Size, antibody dilution of 1:20, apparent MW was 102 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.</p>

Images


Simple Western: VPS34 Antibody [NB110-87320] - Simple Western lane view shows a specific band for VPS34 in 0.5 mg/ml of HepG2 lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.




Western Blot: VPS34 Antibody [NB110-87320] - ATG7 depleted SKBR3 cells show increased sensitivity to ATRA and the RARalpha agonist AM580. RARalpha, ATG5 and VPS34 Western blot analysis of control and the respective SKBR3 knockdown cells. GAPDH was used as a loading control. Image collected and cropped by CiteAb from the following publication (), licensed under a CC-BY license.




Western Blot: VPS34 Antibody [NB110-87320] - HepG2 whole cell lysates.



Immunocytochemistry/Immunofluorescence: VPS34 Antibody [NB110-87320] - VPS34 antibody was tested at 1:200 in HeLa cells with FITC (green). Nuclei (Blue) were counterstained with Dapi (blue).



Western Blot: VPS34 Antibody [NB110-87320] - Celastrol regulates autophagy- & mitophagy-related gene expressions in neurons. SH-SY5Y cells treated with 0.1–1 μ M celastrol for 4 h. (A) Real-time quantitative PCR results show that 1 μ M celastrol treatment for 4 h enhanced mRNA expressions of phagophore induction genes Beclin 1 & Ambra1, phagophore elongation genes Atg7 & Atg12, & vacuole formation genes LC3A & Atg4B. (B) Western blot results of SH-SY5Y treated with 100–500 nM celastrol show celastrol (500 nM) increased in protein expressions of Beclin 1 & Vps34 (phagophore induction), Atg7 (phagophore elongation), LC3-II (vacuole formation), DJ-1, & PINK1 (mitophagy). Celastrol (500 nM) suppressed LRRK2 expression. (C) MPTP (10 mg/kg/day for 3 days, i.p.) but not celastrol (3 mg/kg/day for 3 days, i.p.) caused dopaminergic nerve terminal degeneration (tyrosine hydroxylase \downarrow) & mitophagy inactivation (DJ-1 \downarrow & PINK1 \downarrow) in the striatum of mice as compared to saline control. Celastrol cotreatment with MPTP reversed it as compared to MPTP. Data are given as the mean \pm SEM (n = 3–5). p-value was determined using the Kruskal–Wallis test followed by Dunn's multiple comparison post hoc test. * p < 0.05 compared with controls. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/31906147>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Yang D, Wu X, Wang W et al. Ciliary Type III Adenylyl Cyclase in the VMH Is Crucial for High-Fat Diet-Induced Obesity Mediated by Autophagy *Advanced Science* 2022-01-01 [PMID: 34783461] (Western Blot)

Meena N, Ralston E, Raben N, Puertollano R Enzyme Replacement Therapy Can Reverse Pathogenic Cascade in Pompe Disease *Mol Ther Methods Clin Dev* 2020-07-17 [PMID: 32671132] (FLOW, Human)

Rijal R, Cadena LA, Smith MR et al. Polyphosphate is an extracellular signal that can facilitate bacterial survival in eukaryotic cells *Proceedings of the National Academy of Sciences of the United States of America* 2020-12-15 [PMID: 33268492]

Lin M W, Lin C C et al. Celastrol Inhibits Dopaminergic Neuronal Death of Parkinson's Disease through Activating Mitophagy. *Antioxidants (Basel)* 2019-12-31 [PMID: 31906147] (WB, Mouse)

Campa CC, Margaria JP, Derle A et al. Rab11 activity and PtdIns(3)P turnover removes recycling cargo from endosomes *Nat. Chem. Biol.* 2018-06-18 [PMID: 29915378] (WB)

Brigger D, Schlafli AM, Garattini E, Tschan MP. Activation of RAR α induces autophagy in SKBR3 breast cancer cells and depletion of key autophagy genes enhances ATRA toxicity. *Cell Death Dis* 2015-08-28 [PMID: 26313912] (WB, Human)

Huang Y, Hou JK, Chen TT et al. PML-RAR α enhances constitutive autophagic activity through inhibiting the Akt/mTOR pathway. *Autophagy*. 2011-10-01 [PMID: 21673516] (WB, Human)

Shi JM, Bai LL, Zhang DM et al. Saxifragifolin D induces the interplay between apoptosis and autophagy in breast cancer cells through ROS-dependent endoplasmic reticulum stress. *Biochem Pharmacol* 2013-01-21 [PMID: 23348250] (WB, Human)

Procedures

Serum protocol for VPS34 Antibody (NB110-87320)

VPS34 Antibody:

Western Blot Protocol

1. Perform SDS-PAGE (4-12% MOPS) on samples to be analyzed, loading 40 ug of total protein per lane.
2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
3. Rinse membrane with dH₂O and then stain the blot using Ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
4. Rinse the blot in TBS for approximately 5 minutes.
5. Block the membrane using 5% BSA in TBS + Tween, 1 hour at RT.
6. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
7. Dilute the rabbit anti-VPS34 primary antibody (NB 110-87320) in blocking buffer and incubate 1 hour at room temperature.
8. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
9. Apply the diluted rabbit-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
10. Wash the blot in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturers instructions (Pierce ECL).

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB110-87320

NBL1-14416	VPS34 Overexpression Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB110-87320

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

