

# Product Datasheet

## Survivin [p Ser20] Antibody - BSA Free NB110-92717

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

### Publications: 5

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB110-92717](http://www.novusbio.com/NB110-92717)

Updated 10/23/2024 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB110-92717](http://www.novusbio.com/reviews/destination/NB110-92717)




**NB110-92717**

Survivin [p Ser20] Antibody - BSA Free


Product Information	
Unit Size	0.1 ml
Concentration	1.25 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.1% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS, 30% Glycerol
Target Molecular Weight	16 kDa
Product Description	
Host	Rabbit
Gene ID	332
Gene Symbol	BIRC5
Species	Human, Mouse, Rat
Reactivity Notes	Mouse reactivity reported in scientific literature (PMID: 22814318). Rat reactivity reported in scientific literature (PMID: 17612487). Human reactivity reported in scientific literature (PMID: 24069188).
Immunogen	This Survivin [p Ser20] Antibody was developed against a synthetic peptide surrounding the phosphorylated serine 20 of the human Survivin protein. [Swiss-Prot #O15392]
Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot reported in scientific literature (PMID 24069188), Immunohistochemistry 1:10-1:500. Use reported in scientific literature (PMID 22814318), Immunocytochemistry/ Immunofluorescence 1:100
Application Notes	The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors.

## Images

Western Blot: Survivin [p Ser20] Antibody [NB110-92717] - Detection of Survivin [Ser20] in phosphorylated recombinant protein (+), but not in unphosphorylated protein (-) using [NB110-92717]. Band detected at slightly higher than the predicted molecular weight of 16 kDa.



Immunocytochemistry/Immunofluorescence: Survivin [p Ser20] Antibody [NB110-92717] - Survivin [p Ser20] Antibody (NB110-92717) was tested in HeLa cells with DyLight 488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and DyLight 550 (red).



## Publications

Garlapati C, Joshi S, Bhattarai S et al. PLK1 and AURKB phosphorylate survivin differentially to affect proliferation in racially distinct triple-negative breast cancer Cell Death & Disease 2023-01-10 [PMID: 36627281] (Immunohistochemistry)

Benaiges E, Ceperuelo-Mallafre V, Madeira A et al. Survivin drives tumor-associated macrophage reprogramming: a novel mechanism with potential impact for obesity Cellular oncology (Dordrecht) 2021-03-12 [PMID: 33710603]

Knapp Charles F, Sayegh Zena, Schell Michael J et al. Expression of CXCR4, E-cadherin, Bcl-2, and survivin in Merkel cell carcinoma: an immunohistochemical study using a tissue microarray. The American Journal of Dermatopathology 2012-01-01 [PMID: 22814318] (IF/IHC, Mouse)

Muschol-Steinmetz C, Friemel A, Kreis NN et al. Function of Survivin in Trophoblastic Cells of the Placenta. PLoS One. 2013-09-19 [PMID: 24069188] (WB, Human)

Dohi, T et al. Compartmentalized phosphorylation of IAP by protein kinase A regulates cytoprotection. Mol Cell;27 (1):17-28. 2007-07-06 [PMID: 17612487] (WB, Rat)

## Procedures

### **Immunocytochemistry/Immunofluorescence Protocol for Survivin [p Ser20] antibody (NB110-92717)**

Survivin [p Ser20] Antibody:

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.
4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,000 and incubate for 10 minutes. Wash a third time for 10 minutes.
9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

\*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NB110-92717**

---

NBL1-07987	Survivin Overexpression Lysate
NB110-92717PEP	Survivin Antibody Blocking Peptide
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB110-92717](http://www.novusbio.com/reviews/submit/NB110-92717)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

