Product Datasheet

CHRND Antibody (88B) NB120-2804

Unit Size: 100uL

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB120-2804

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB120-2804



NB120-2804

CHRND Antibody (88B)

100uL
This product is unpurified. The exact concentration of antibody is not quantifiable.
Store at -20C. Avoid freeze-thaw cycles.
Monoclonal
88B
0.05% Sodium Azide
IgG1
Unpurified
Ascites
Mouse
1144
CHRND
Human, Mouse, Rat, Amphibian, Chicken, Fish
Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Additional Mouse on Mouse blocking steps may be required for IHC and ICC experiments. Please contact Technical Support for more information.
Detects nicotinic acetylcholine receptor (nAChR) gamma and delta subunits in torpedo and the delta subunit in mouse, rat, frog and chicken. Does not detect the alpha 1 and beta 1 subunits.
Purified Torpedo californica acetylcholine receptor.
Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry- Paraffin, Immunoprecipitation
Western Blot 1:5000, Immunohistochemistry 1:1000, Immunocytochemistry/ Immunofluorescence 1:20 - 100, Immunoprecipitation 1:10 - 1:500, Immunohistochemistry-Paraffin 1:1000, Immunohistochemistry-Frozen 1:1000
WB: Detects an approx. 60 kDa protein representing the gamma subunit and a 65 kDa protein representing the delta subunit from torpedo skeletal muscle homogenates. Under non-reducing conditions the delta subunit migrates mostly as a dimer of ~130 kDa. IHC: Staining of nAChR in rat skeletal muscle results in strong staining of the motor endplate. Cryostat or permeabilization of the sections is recommended due to the reactivity of this antibody with the cytoplasmic side of the receptor.

www.novusbio.com



×


Images

Immunocytochemistry/Immunofluorescence: CHRND Antibody (88B) [NB120-2804] - Analysis of Nicotinic Acetylcholine Receptor using Anti-Nicotinic Acetylcholine Receptor Monoclonal Antibody (88B) shows staining in U251 Cells. Nicotinic Acetylcholine Receptor staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Nicotinic Acetylcholine Receptor at a dilution of 1:20 over night at 4C, washed with PBS and incubated with a DyLight-488 conjugated.

Immunohistochemistry-Paraffin: CHRND Antibody (88B) [NB120-2804] - Normal biopsies of deparaffinized Mouse lymph node tissue.


Immunocytochemistry/Immunofluorescence: CHRND Antibody (88B) [NB120-2804] - Analysis of Nicotinic Acetylcholine Receptor using Anti-Nicotinic Acetylcholine Receptor Monoclonal Antibody (88B) shows staining in Hela Cells. Nicotinic Acetylcholine Receptor staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Nicotinic Acetylcholine Receptor at a dilution of 1:100 over night at 4C, washed with PBS and incubated with a DyLight-488 conjugated.

Immunocytochemistry/Immunofluorescence: CHRND Antibody (88B) [NB120-2804] - Analysis of Nicotinic Acetylcholine Receptor using Anti-Nicotinic Acetylcholine Receptor Monoclonal Antibody (88B) shows staining in Neuro-2a Cells. Nicotinic Acetylcholine Receptor staining (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were probed without (control) or with or an antibody recognizing Nicotinic Acetylcholine Receptor at a dilution of 1:100 over night at 4C, washed with PBS and incubated with a DyLight-488 conjugated.






Immunohistochemistry-Paraffin: CHRND Antibody (88B) [NB120-2804] - Normal biopsies of deparaffinized Mouse brain tissue.



Immunohistochemistry-Paraffin: CHRND Antibody (88B) [NB120-2804] -Normal biopsies of deparaffinized Mouse kidney tissue.



Publications

Li B, Chen L, Gu Yd Stability of motor endplates is greater in the biceps than in the interossei in a rat model of obstetric brachial plexus palsy Neural Regen Res 2020-09-01 [PMID: 32209772] (WB, Rat)

Dmytrenko GM, Pumplin DW, Bloch RJ. Dystrophin in a membrane skeletal network: localization and comparison to other proteins. J Neurosci;13(2):547-58. 1993-02-01 [PMID: 8426227]

Bloch RJ, Sealock R, Pumplin DW, Luther PW, Froehner SC. Association of acetylcholine receptors with peripheral membrane proteins: evidence from antibody-induced coaggregation. J Membr Biol;138(1):13-28. 1994-02-01 [PMID: 8189428]

Froehner SC, Douville K, Klink S, Culp WJ. Monoclonal antibodies to cytoplasmic domains of the acetylcholine receptor. J Biol Chem;258(11):7112-20. 1983-06-10 [PMID: 6189834]







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NB120-2804

H00001144-Q01-10ug	Recombinant Human CHRND GST (N-Term) Protein
NBP1-97005-0.5mg	Mouse IgG1 Isotype Control (MG1)
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB120-2804

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

