

Product Datasheet

GAPDH Antibody NB300-325

Unit Size: 0.1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 2 Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB300-325

Updated 4/13/2025 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB300-325



NB300-325**GAPDH Antibody****Product Information**

Unit Size	0.1 ml
Concentration	0.2 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	TBS and 0.1% BSA
Target Molecular Weight	36 kDa

Product Description

Host	Rabbit
Gene ID	2597
Gene Symbol	GAPDH
Species	Human, Mouse, Rat
Reactivity Notes	Human and mouse.
Marker	Cytosolic Marker
Immunogen	This GAPDH antibody was developed against an epitope between residue 300 and the C-terminus of the human GAPDH protein [accession number NP_002037.2]


Product Application Details

Applications	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Immunoprecipitation
Recommended Dilutions	Western Blot 1:2000-1:10000, Simple Western 1:500, Immunocytochemistry/ Immunofluorescence 1:250, Immunoprecipitation
Application Notes	<p>This GAPDH antibody is useful for Immunocytochemistry/Immunofluorescence and Western blot, where a band is observed approx. 36 kDa.</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.</p> <p>See Simple Western Antibody Database for Simple Western validation: Tested in HeLa lysate 0.05 mg/mL, separated by Size, antibody dilution of 1:500, apparent MW was 42 kDa. Separated by Size-Wes, Sally Sue/Peggy Sue.</p> <p>The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors.</p>




Images


Western Blot: GAPDH Antibody [NB300-325] - Detection of mouse and rat GAPDH by western blot. Samples: Whole cell lysate (50 ug) from NIH 3T3, TCMK-1, 4T1, CT26.WT, and rat C6 cells. Antibodies: Affinity purified rabbit anti-GAPDH antibody NB300-325 used for WB at 0.5 ug/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.




Immunocytochemistry/Immunofluorescence: GAPDH Antibody [NB300-325] - GAPDH antibody was tested in HeLa cells with DyLight 488 (green). Nuclei and alpha-tubulin were counterstained with DAPI (blue) and DyLight 550 (red).



Western Blot: GAPDH Antibody [NB300-325] - Whole cell lysate (50 ug) from HeLa (H) and 293T (50 ug) probed with GAPDH Antibody diluted at 0.04 ug/ml




Western Blot: GAPDH Antibody [NB300-325] - Detection of Mouse GAPDH Whole cell lysate (5, 15 and 50 ug) from mouse NIH3T3 cells. Another Affinity purified rabbit anti-GAPDH used at 0.04 mcg/ml. Chemiluminescence with an exposure time of 30 seconds.




Simple Western: GAPDH Antibody [NB300-325] - Simple Western lane view shows a specific band for GAPDH in 0.05 mg/ml of HeLa lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.




Western Blot: GAPDH Antibody [NB300-325] - GCNY-mediated serum decreased Nrf2 levels & inflammatory cytokines of NF- κ B & IL-6. (A) Intracellular relative ROS levels in HMC-3 cells treated with different concentrations (0, 50, 100, & 200 μ M) of H₂O₂ for 4 h. Auto-fluo refers to the auto-fluorescence intensity value of HMC-3 cells. (B,C) The mRNA & protein levels of Nrf2 were expressed by HMC-3 treated with different concentrations of H₂O₂. (D) The significantly elevated whole protein level in NC + H₂O₂ group cells indicates an increased proliferation of HMC-3 cells compared to the NC group cells & (E) ROS levels were attenuated in cells treated by GCNY-mediated serum. (F-H) Through automated Western blot analysis, GCNY group cells expressed lower levels of Nrf2, NF- κ B p65, pNF- κ B p65 (ser536), & IL-6 compared with NC + H₂O₂ group cells. ns, no significance, * p < 0.05; ** p < 0.01; *** p < 0.001; **** p < 0.0001. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/36188472>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.




Western Blot: GAPDH Antibody [NB300-325] - GCNY-mediated serum decreased Nrf2 levels & inflammatory cytokines of NF- κ B & IL-6. (A) Intracellular relative ROS levels in HMC-3 cells treated with different concentrations (0, 50, 100, & 200 μ M) of H₂O₂ for 4 h. Auto-fluo refers to the auto-fluorescence intensity value of HMC-3 cells. (B,C) The mRNA & protein levels of Nrf2 were expressed by HMC-3 treated with different concentrations of H₂O₂. (D) The significantly elevated whole protein level in NC + H₂O₂ group cells indicates an increased proliferation of HMC-3 cells compared to the NC group cells & (E) ROS levels were attenuated in cells treated by GCNY-mediated serum. (F-H) Through automated Western blot analysis, GCNY group cells expressed lower levels of Nrf2, NF- κ B p65, pNF- κ B p65 (ser536), & IL-6 compared with NC + H₂O₂ group cells. ns, no significance, * p < 0.05; ** p < 0.01; *** p < 0.001; **** p < 0.0001. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/36188472>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: GAPDH Antibody [NB300-325] - GCNY-mediated serum decreased Nrf2 levels & inflammatory cytokines of NF- κ B & IL-6. (A) Intracellular relative ROS levels in HMC-3 cells treated with different concentrations (0, 50, 100, & 200 μ M) of H₂O₂ for 4 h. Auto-fluo refers to the auto-fluorescence intensity value of HMC-3 cells. (B,C) The mRNA & protein levels of Nrf2 were expressed by HMC-3 treated with different concentrations of H₂O₂. (D) The significantly elevated whole protein level in NC + H₂O₂ group cells indicates an increased proliferation of HMC-3 cells compared to the NC group cells & (E) ROS levels were attenuated in cells treated by GCNY-mediated serum. (F–H) Through automated Western blot analysis, GCNY group cells expressed lower levels of Nrf2, NF- κ B p65, pNF- κ B p65 (ser536), & IL-6 compared with NC + H₂O₂ group cells. ns, no significance, * p < 0.05; ** p < 0.01; *** p < 0.001; **** p < 0.0001. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/36188472>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Western Blot: GAPDH Antibody [NB300-325] - GCNY-mediated serum decreased Nrf2 levels & inflammatory cytokines of NF- κ B & IL-6. (A) Intracellular relative ROS levels in HMC-3 cells treated with different concentrations (0, 50, 100, & 200 μ M) of H₂O₂ for 4 h. Auto-fluo refers to the auto-fluorescence intensity value of HMC-3 cells. (B,C) The mRNA & protein levels of Nrf2 were expressed by HMC-3 treated with different concentrations of H₂O₂. (D) The significantly elevated whole protein level in NC + H₂O₂ group cells indicates an increased proliferation of HMC-3 cells compared to the NC group cells & (E) ROS levels were attenuated in cells treated by GCNY-mediated serum. (F–H) Through automated Western blot analysis, GCNY group cells expressed lower levels of Nrf2, NF- κ B p65, pNF- κ B p65 (ser536), & IL-6 compared with NC + H₂O₂ group cells. ns, no significance, * p < 0.05; ** p < 0.01; *** p < 0.001; **** p < 0.0001. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/36188472>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Chen Y, Wang B, Lai WF et al. Chinese herbal formula (GCNY)-mediated serum alleviates peroxidation induced by H₂O₂ in human microglial cells *Frontiers in Neuroscience* 2022-09-14 [PMID: 36188472] (Western Blot, Block/Neutralize)

Brinks J, van Dijk EHC, Kielbasa SM et al. The Cortisol Response of Male and Female Choroidal Endothelial Cells: Implications for Central Serous Chorioretinopathy *The Journal of clinical endocrinology and metabolism* 2021-09-21 [PMID: 34546342] (WB, Human)

Ahmed AA, Neidle S A G-Quadruplex-Binding Small Molecule and the HDAC Inhibitor SAHA (Vorinostat) Act Synergistically in Gemcitabine-Sensitive and Resistant Pancreatic Cancer Cells *Sci Adv* 2020-11-01 [PMID: 33227941] (WB, Human)

Rosen MB, Jeffay SC, Nichols HP et al. ATP Binding Cassette Sub-family Member 2 (ABCG2) and Xenobiotic Exposure During Early Mouse Embryonic Stem Cell Differentiation. *Birth Defects Res.* 2017-10-09 [PMID: 28990372] (WB, Mouse)

Procedures

Western Blot protocol for GAPDH Antibody (NB300-325)

GAPDH Antibody:

Western Blot Protocol

1. Perform SDS-PAGE (4-12% MOPS) on samples to be analyzed, loading 25 ug of total protein per lane.
 2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
 3. Rinse membrane with dH₂O and then stain the blot using Ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
 4. Rinse the blot in TBS for approximately 5 minutes.
 5. Block the membrane using 5% NFDM + 1% BSA in TBS + Tween, 1 hour at RT.
 6. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
 7. Dilute the rabbit anti-GAPDH primary antibody (NB300-325) in blocking buffer and incubate 1 hour at room temperature.
 8. Rinse the membrane in dH₂O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
 9. Apply the diluted rabbit-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
 10. Wash the blot in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each (this step can be repeated as required to reduce background).
 11. Apply the detection reagent of choice in accordance with the manufacturers instructions (Pierce ECL).
- Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.

Immunocytochemistry/Immunofluorescence protocol for GAPDH Antibody (NB300-325)

GAPDH Antibody:

Culture cells to appropriate density in 35 mm culture dishes or 6-well plates.

1. Remove culture medium and add 10% formalin to the dish. Fix at room temperature for 30 minutes.
2. Remove the formalin and add ice cold methanol. Incubate for 5-10 minutes.
3. Remove methanol and add washing solution (i.e. PBS). Be sure to not let the specimen dry out. Wash three times for 10 minutes.
4. To block nonspecific antibody binding incubate in 10% normal goat serum from 1 hour to overnight at room temperature.
5. Add primary antibody at appropriate dilution and incubate at room temperature from 2 hours to overnight at room temperature.
6. Remove primary antibody and replace with washing solution. Wash three times for 10 minutes.
7. Add secondary antibody at appropriate dilution. Incubate for 1 hour at room temperature.
8. Remove antibody and replace with wash solution, then wash for 10 minutes. Add Hoechst 33258 to wash solution at 1:25,000 and incubate for 10 minutes. Wash a third time for 10 minutes.
9. Cells can be viewed directly after washing. The plates can also be stored in PBS containing Azide covered in Parafilm (TM). Cells can also be cover-slipped using Fluoromount, with appropriate sealing.

*The above information is only intended as a guide. The researcher should determine what protocol best meets their needs. Please follow safe laboratory procedures.



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB300-325

NBL1-10967	GAPDH Overexpression Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB300-325

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

