

# Product Datasheet

## PDE10A Antibody Blocking Peptide NB300-645PEP

Unit Size: 0.25 mg

Store at -20C. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NB300-645PEP](http://www.novusbio.com/NB300-645PEP)

Updated 10/23/2024 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NB300-645PEP](http://www.novusbio.com/reviews/destination/NB300-645PEP)



**NB300-645PEP****PDE10A Antibody Blocking Peptide**

<b>Product Information</b>	
<b>Unit Size</b>	0.25 mg
<b>Concentration</b>	2.5 mg/ml
<b>Storage</b>	Store at -20C. Avoid freeze-thaw cycles.
<b>Purity</b>	99.9
<b>Buffer</b>	Sodium Bicarbonate buffer, pH 8.00 containing up to 10% DMSO
<b>Product Description</b>	
<b>Description</b>	Synthetic peptide (Human) taken against a common sequence near the 250-325 amino acids downstream of N terminal region of PDE10A. Synthetic MAP. GenBank Accession#: AAD32596.1
<b>Gene ID</b>	10846
<b>Gene Symbol</b>	PDE10A
<b>Species</b>	Human
<b>Product Application Details</b>	
<b>Applications</b>	Antibody Competition
<b>Recommended Dilutions</b>	Antibody Competition
<b>Application Notes</b>	This peptide is useful as a blocking peptide for NB300-645. For further blocking peptide related protocol, click <a href="#">here</a> .



## Procedures

### Antibody Competition protocol for PDE10A Protein (NB300-645PEP)

Antibody Competition Protocol for PDE10A Protein (NB300-645PEP):

In this assay the antigen binding sites (Fab) of a particular antibody is allowed to bind to homologous antigenic peptide in several hundred molar excess ratio of immunoglobulin to peptide. Since the antigenic peptide has higher affinity for Fab site than the full length native proteins, the antigen binding site is blocked thus antibodies are unable to bind to the specific sites on either native, partial denatured or completely denature antigen on sections, in solution or on membrane blots.

Before starting, standardize the conditions for Western blot, immunoprecipitation or immunohistochemistry using the relevant antibody. Conditions include: volume of antibody used; dilution factor, final volume, incubation time, washing conditions etc. Once conditions have been standardized, repeat the same protocol in duplicate using blocked and unblocked antibody.

Begin the competition assay by preparing the blocked antibody-peptide solution:

1. Take the same volume of antibody as previously standardized. Let us assume 10 uL in 15 mL.
2. Add approximately 1:200 moles of excess peptide (peptides are generally 2200 dalton). This will be equivalent of approximately 60-70ul of peptide solution (original peptide concentration is 2.5mg/ml).
3. Mix 10 uL of antibody with 60-70 uL of peptide and make up the volume to 200 uL using DiluOBuffer.

Compare the above blocked antibody-peptide solution with control antibody:

1. Take same amount of antibody as used in the blocked antibody-peptide solution and add 60-70 ul of PBS and make up the volume to 200 ul using DiluOBuffer.
2. Once both blocked antibody-peptide solution and the control antibody solution have been made, incubate both solutions at 4 degrees C overnight on a rotating mixer. Centrifuge both tubes at 12,000-14000 xg for 1-2 minute at 4 degrees C. A small amount of precipitate may accumulate in the blocked antibody-peptide solution due to antibody-antigen complex formation. If this occurs, carefully remove the precipitate and use only the supernatant.
3. Dilute both solutions in 1X DiluOBuffer to the final dilution volume of 15 mL. The antibody-peptide solution will not give any labeling or will have significantly reduced labeling while the Control Antibody solution should provide standard results.



**Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

**Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

**Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

**General Contact Information**

[www.novusbio.com](http://www.novusbio.com)  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

**Products Related to NB300-645PEP**

---

H00010846-Q01-10ug	Recombinant Human PDE10A GST (N-Term) Protein
AF467	EphB2 Antibody [Unconjugated]
NB300-645-0.05mg	PDE10A Antibody
273-F9-025	FGF-9 [Unconjugated]

---

**Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Peptides and proteins are guaranteed for 3 months from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NB300-645PEP](http://www.novusbio.com/reviews/submit/NB300-645PEP)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

