

Product Datasheet

p38 alpha [p Thr180, p Tyr182] Antibody - Azide Free NB500-138

Unit Size: 0.1 ml

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Publications: 17

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB500-138

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB500-138



NB500-138


p38 alpha [p Thr180, p Tyr182] Antibody - Azide Free

Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	No Preservative
Isotype	IgG
Purity	Antigen Affinity-purified
Buffer	10mM HEPES (pH 7.5), 0.15M NaCl, 0.1 mg/ml BSA and 50% Glycerol
Target Molecular Weight	39 kDa
Product Description	
Host	Rabbit
Gene ID	1432
Gene Symbol	MAPK14
Species	Human, Mouse, Zebrafish
Reactivity Notes	Zebrafish reactivity reported in scientific literature (PMID: 29692779). Mouse reactivity reported in scientific literature (PMID: 24105651).
Specificity/Sensitivity	Specific for endogenous levels of the ~39 kDa p38 alpha protein phosphorylated at Thr180/Tyr182. Immunolabeling is blocked by preadsorption with the phosphopeptide used as antigen, but not by the corresponding non-phosphopeptide.
Immunogen	Synthetic phospho-peptide corresponding to amino acid residues surrounding Thr180/Tyr182 conjugated to KLH. Accession # P70618
Product Application Details	
Applications	Western Blot, Flow (Intracellular), Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1:1000, Immunohistochemistry 1:250, Immunohistochemistry-Paraffin 1:250, Flow (Intracellular)
Application Notes	Use in Flow-intracellular reported in scientific literature (PMID: 24105651).




Images


Western Blot: p38 [p Thr180, p Tyr182] Antibody [NB500-138] - HeLa cell lysates that had been treated with UV or untreated (Control) showing specific immunolabeling of the ~39 kDa p38 MAPK protein phosphorylated at Thr180/Tyr182.




Immunohistochemistry-Paraffin: p38 [p Thr180, p Tyr182] Antibody [NB500-138] - Immunostaining of human breast cancer tissue showing p38 when phosphorylated at Thr180/Tyr182 in red.



Western Blot: p38 [p Thr180, p Tyr182] Antibody [NB500-138] - Western blot of HeLa cell lysates that had been treated with UV or untreated (Control) showing specific immunolabeling of the ~39k p38 MAPK protein phosphorylated at Thr180/Tyr182.



Western Blot: p38 alpha [p Thr180, p Tyr182] Antibody [NB500-138] - RIP2 is not essential for MAPK pathways in the early ontogenesis. (A) Immunoblot analysis of phospho-p44/42 MAPK, p44/42 MAPK, & p38 MAPK in larvae homogenate from WT & NOD1-1IS^{-/-} zebrafish at 7 days post-fertilization (dpf). (B) Immunoblot analysis of phospho-p44/42 MAPK, p44/42 MAPK, & p38 MAPK in larvae homogenate from WT & RIP2^{-/-} zebrafish at 7 dpf. (C) Immunoblot analysis of Atg5, p62, & LC3b in larvae homogenate from WT & NOD1-1IS^{-/-} zebrafish at 7 dpf. (D) Immunoblot analysis of Atg5, p62, & LC3b in larvae homogenate from WT & RIP2^{-/-} zebrafish at 7 dpf. Western blotting results were quantified using Quantity One software. Data represent the average of two independent experiments. *p < 0.05, **p < 0.01. Image collected & cropped by CiteAb from the following publication (<https://pubmed.ncbi.nlm.nih.gov/29692779>), licensed under a CC-BY license. Not internally tested by Novus Biologicals.



Publications

Ren J Phosvitin Extraction from Egg Yolk and Its Potential as a Functional Food Ingredient for Improving Bone Health J Sci Food Agric 2015-01-08 [PMID: 25565664]

Mukherjee U, Samanta A, Biswas S Et al. Chronic exposure to nonylphenol induces oxidative stress and liver damage in male zebrafish (Danio rerio): Mechanistic insight into cellular energy sensors, lipid accumulation and immune modulation Chemico-biological interactions 2021-11-26 [PMID: 34843692]

Zhao S, Xu J, Zhang W et al. Paternal exposure to microcystin-LR triggers developmental neurotoxicity in zebrafish offspring via an epigenetic mechanism involving MAPK pathway The Science of the total environment 2021-06-16 [PMID: 34153754]

Ren J, Chakrabarti S, Wu J Inhibition of osteoclastogenesis and inflammation by phosvitin and phosvitin hydrolysate via NF-kappa B and MAPK pathways in RAW 264.7 cells Journal of Food Bioactives 2021-03-31

Sirin S, Aslim B Protective effect of exopolysaccharides from lactic acid bacteria against amyloid beta1-42induced oxidative stress in SH-SY5Y cells: Involvement of the AKT, MAPK, and NF- kappa B signaling pathway Process Biochemistry Jul 1 2021 12:00AM

Zhao S, Yuan C, Tuo X et al. MCLR induces dysregulation of calcium homeostasis and endoplasmic reticulum stress resulting in apoptosis in Sertoli cells Chemosphere 2020-08-15 [PMID: 32828052]

Mukherjee U, Samanta A, Biswas S et al. Bisphenol A-induced oxidative stress, hepatotoxicity and altered estrogen receptor expression in Labeo bata: impact on metabolic homeostasis and inflammatory response Ecotoxicology and Environmental Safety 2020-10-01 [PMID: 32800225] (WB, Mouse)

Zhang Y, Beketaev I, Segura AM et al. Contribution of Increased Expression of Yin Yang 2 to Development of Cardiomyopathy Front Mol Biosci 2020-03-03 [PMID: 32195266] (WB, Mouse)

Singh A, Jones OD, Mockett BG et al. Tumor Necrosis Factor-alpha-Mediated Metaplastic Inhibition of LTP is Constitutively Engaged in an Alzheimer's Disease Model J. Neurosci. 2019-09-30 [PMID: 31570539] (WB, Mouse, Rat)

Ren J Phosvitin Extraction from Egg Yolk and Its Potential as a Functional Food Ingredient for Improving Bone Health Thesis (WB)

Pal S, Nath P, Biswas S et al. Nonylphenol attenuates SOCS3 expression and M1 polarization in lipopolysaccharide-treated rat splenic macrophages Ecotoxicol. Environ. Saf. 2019-06-15 [PMID: 30870658] (WB, Rat)

Ohm AM, Affandi T, Reyland ME. EGF receptor and PKCd kinase activate DNA damage-induced pro-survival and pro-apoptotic signaling via biphasic activation of ERK and MSK1 kinases. J. Biol. Chem. 2019-01-24 [PMID: 30679314] (WB, Rat)

More publications at <http://www.novusbio.com/NB500-138>



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB500-138

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NB100-56665PEP	p38 alpha Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB500-138

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

