# **Product Datasheet**

# ICAM-1/CD54 Antibody (MEM-111) - BSA Free NB500-318

Unit Size: 0.1 mg

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

**Publications: 7** 

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB500-318

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB500-318



#### NB500-318

ICAM-1/CD54 Antibody (MEM-111) - BSA Free

| • (                        | ,                           |
|----------------------------|-----------------------------|
| <b>Product Information</b> |                             |
| Unit Size                  | 0.1 mg                      |
| Concentration              | 1 mg/ml                     |
| Storage                    | Store at 4C. Do not freeze. |
| Clonality                  | Monoclonal                  |
| Clone                      | MEM-111                     |
| Preservative               | 0.9% Sodium Azide           |
| Isotype                    | IgG2a                       |
| Purity                     | Protein A purified          |
| Buffer                     | PBS (pH 7.4)                |
| Target Molecular Weight    | 110 kDa                     |
| Product Description        |                             |
| Host                       | Mouse                       |
| Gene ID                    | 3383                        |


| Gene ID                 | 3383  |
|-------------------------|---|
| Gene Symbol             | ICAM1   |
| Species                 | Human, Rat, Bovine  |
| Specificity/Sensitivity | The antibody MEM-111 reacts with an extracellular epitope CD54 (ICAM-1), a 85 -110 kDa type I transmembrane glycoprotein (receptor for rhinovirus). The expression of CD54 is upregulated by activation; it is expressed on activated endothelial cells, T lymphocytes, B lymphocytes, monocytes, macrophages, granulocytes and dendritic cells. HLDA VI; WS Code AS A049 |
| Immunogen               | Raji human Burkitt's lymphoma cell line (NM_000201.2).  |
|                         |   |

| Product Application Details |   |
|-----------------------------|---|
| Applications                | Western Blot, ELISA, Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Frozen, Immunohistochemistry-Paraffin, CyTOF-ready  |
| Recommended Dilutions       | Western Blot 1-2 ug/ml, Flow Cytometry 0.5-4 ug/ml, ELISA 1:2000, Immunohistochemistry 1:10 - 1:500, Immunohistochemistry-Paraffin 10 ug/ml, Immunohistochemistry-Frozen, CyTOF-ready |
| Application Notes           | WB: Use under non-reducing conditions. Use in IHC-Frozen reported in scientific literature (PMID 26256764). This antibody is CyTOF ready.   |

#### **Images**

license.

Western Blot: ICAM-1/CD54 Antibody (MEM-111) [NB500-318] - ICAM-1 protein expression levels in rat lung tissue after the different treatments in each group. Immunoblotting showed increase in ICAM-1 after one course of paclitaxel compared with normal lung tissue; p < 0.01. These changes were reversed by using parecoxib in combination with paclitaxel, while the group that received parecoxib treatment alone resembled the Con group. The bar graph under the protein strap represents the corresponding band intensities; the data are presented as x + - s and corrected by GA. Image collected and cropped by CiteAb from the following publication (https://www.nature.com/articles/srep12977), licensed under a CC-BY





Flow Cytometry: ICAM-1/CD54 Antibody (MEM-111) [NB500-318] - Flow cytometry analysis (surface staining) of CD54 expression in activated human peripheral blood with anti-CD54 (MEM-111) PE SS-H :: SS INT LIN CD<sub>54</sub> PE Western Blot: ICAM-1/CD54 Antibody (MEM-111) [NB500-318] - Of kDa: B CD54 expression in activated (A) and nonactivated (B) Huvec cells by 103 antibody MEM-111. Lower bands represent tubulin as a loading control. 81 37 Flow Cytometry: ICAM-1/CD54 Antibody (MEM-111) [NB500-318] -Analysis using the FITC conjugate of NB500-318. Staining of Human peripheral blood lymphocytes with MEM-111 FITC. 102 Log Fluoresence Intensity Flow Cytometry: ICAM-1/CD54 Antibody (MEM-111) [NB500-318] -Analysis using PE conjugate of NB500-318. A cell surface stain was 300 performed on THP-1 cells with ICAM-1/CD54 antibody (MEM-111) Relative Cell Number NB500-318 (blue) and a matched isotype control (orange). Cells were incubated in an antibody dilution of 1:100 for 20 minutes at room 200 temperature. Both antibodies were conjugated to PE. 100 102 ICAM-1/CD54 Antibody (MEM-111) [PE]



#### **Publications**

Chung NPY, Khan KMF, Kaner RJ et al. HIV induces airway basal progenitor cells to adopt an inflammatory phenotype Scientific reports 2021-02-17 [PMID: 33597552] (WB, Human)

Carlos CP, Silva AA, Gil CD, Oliani SM. Pharmacological treatment with galectin-1 protects against renal ischaemia-reperfusion injury Sci Rep 2018-06-22 [PMID: 29934646] (IHC-P, Rat)

Thomas A, Ou-Yang HD, Lowe-Krentz L. Biomimetic channel modeling local vascular dynamics of pro-inflammatory endothelial changes. Biomicrofluidics. 2016-01-06 [PMID: 26858813]

Liu WJ, Zhong ZJ, Cao LH et al. Paclitaxel-induced lung injury and its amelioration by parecoxib sodium. Sci Rep. 2015-08-10 [PMID: 26256764] (IHC-Fr, Rat)

Xiang S, Dong NG, Liu JP et al. Inhibitory effects of suppressor of cytokine signaling 3 on inflammatory cytokine expression and migration and proliferation of IL-6/IFN-gamma-induced vascular smooth muscle cells. J Huazhong Univ Sci Technolog Med Sci. 2013-10-01 [PMID: 24142708] (WB, Rat)

Rosano JM, Tousi N, Scott RC et al. A physiologically realistic in vitro model of microvascular networks. Biomed Microdevices 2009-10-01 [PMID: 19452279] (Bovine)

Ng CY, Kamisah Y, Faizah O, Jaarin K. The role of repeatedly heated soybean oil in the development of hypertension in rats: association with vascular inflammation Int J Exp Pathol 2012-10-01 [PMID: 22974219] (IF/IHC, Rat)





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### **Products Related to NB500-318**

NBL1-11804 ICAM-1/CD54 Overexpression Lysate

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-96778 Mouse IgG2a Isotype Control (M2A)

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB500-318

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

