Product Datasheet

VEGFR1/Flt-1 Antibody (EWC) - Azide and BSA Free NB600-1004

Unit Size: 0.1 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.



Reviews: 3 Publications: 2

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB600-1004

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB600-1004



NB600-1004

VEGFR1/Flt-1 Antibody (EWC) - Azide and BSA Free

| Product Information | |
|-----------------------------|---|
| Unit Size | 0.1 mg |
| Concentration | LYOPH mg/ml |
| Storage | Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles. |
| Clonality | Monoclonal |
| Clone | EWC |
| Preservative | No Preservative |
| Reconstitution Instructions | Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml. Please note the sample size of this product will be provided in reconstituted liquid form. |
| Isotype | IgG1 |
| Purity | Protein G purified |
| Buffer | PBS |
| Product Description | |
| Host | Mouse |
| Gene ID | 2321 |
| Gene Symbol | FLT1 |
| Species | Human, Mouse |
| Reactivity Notes | Please note that this antibody is reactive to Mouse and derived from the same host, Mouse. Mouse-On-Mouse blocking reagent may be needed for IHC and ICC experiments to reduce high background signal. You can find these reagents under catalog numbers PK-2200-NB and MP-2400-NB. Please contact Technical Support if you have any questions. |
| Marker | Endothelial Cell Marker |
| Specificity/Sensitivity | The monoclonal will detect native and denaturated human VEGFR-1/ Flt-1 in ELISA experiments and on the surface of different human cell types. The will also detect mouse Flt-1. |
| Immunogen | Recombinant human soluble extracellular Flt-1 (D5) |
| Product Application Details | |
| Applications | Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, In vivo assay |
| Recommended Dilutions | Western Blot 2-5 ug/ml, ELISA 1-10 ug/ml, Immunocytochemistry/ Immunofluorescence 2-10 ug/ml, In vivo assay |
| Application Notes | Not tested in other applications. Optimal dilutions/concentrations should be determined by the end user. Use In vivo reported in scientific literature (PMID:31877123) |

www.novusbio.com



Images

mattaone Western Blot: VEGFR1/Flt-1 Antibody (Flt-1/EWC) [NB600-1004] -Analysis of recombinant human and mouse soluble VEGF receptors using a monoclonal antibody directed against human recombinant sFlt-1 (D5). There is a cross reactivity with mouse sFIt-1(D7)-Fc but no with 170 human and mouse sKDR and human sFLT-4 visible. 100 70 55 40 35 (anti-human VEGFR-1 [101-M24]; SDS-PAGE 10%, red.) Immunocytochemistry/Immunofluorescence: VEGFR1/Flt-1 Antibody (EWC) - Azide and BSA Free [NB600-1004] - VEGFR1/Flt-1 Antibody (Flt-1/EWC) [NB600-1004] - Staining of human Flt-1 in a co-culture of PAE-Flt-1, PAE-KDR and PAE-FLT-4 with a mouse anti-human Flt-1 antibody. Conjugated secondary antibody: goat anti-mouse ALEXA Flour (1:600).Western Blot: VEGFR1/Flt-1 Antibody (Flt-1/EWC) [NB600-1004] -Analysis of recombinant mouse soluble VEGFR-1 conditioned supernatant using a monoclonal antibody directed against human recombinant sFlt-1(D5). There is a strong cross reactivity with the mouse 170 sFlt-1 visible. 100 70 55 40 35 (anti-human Flt-1 [101-M24]; SDS.PAGE 10% red)

Publications

Verma M, Shimizu-Motohashi Y, Asakura Y et al. Inhibition of FLT1 ameliorates muscular dystrophy phenotype by increased vasculature in a mouse model of Duchenne muscular dystrophy PLoS Genet. 2019-12-01 [PMID: 31877123] (In Vivo, Mouse)

Ptaszynska MM, Pendrak ML, Bandle RW, Stracke ML, Roberts DD. Positive feedback between vascular endothelial growth factor-A and autotaxin in ovarian cancer cells. Mol Cancer Res;6(3):352-63. 2008-03-01 [PMID: 18337445]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NB600-1004

| NB100-527PEP | VEGFR1/Flt-1 Antibody Blocking Peptide |
|------------------|---|
| NBP1-97005-0.5mg | Mouse IgG1 Isotype Control (MG1) |
| | |
| NB720-B | Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin] |
| HAF007 | Goat anti-Mouse IgG Secondary Antibody [HRP] |

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB600-1004

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

