Product Datasheet

KMT2A/MLL Antibody - BSA Free NB600-256

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Reviews: 2 Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB600-256

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NB600-256



NB600-256

KMT2A/MLL Antibody - BSA Free


KMT2A/MLL Antibody - BSA Free	
Product Information	
Unit Size	100 ul
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)
Product Description	
Description	Novus Biologicals Rabbit KMT2A/MLL Antibody - BSA Free (NB600-256) is a polyclonal antibody validated for use in WB, Flow, Simple Western, IP and ChIP. Anti-KMT2A/MLL Antibody: Cited in 4 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	4297
Gene Symbol	KMT2A
Species	Human
Immunogen	The immunogen recognized by this antibody maps to a region between residues 2725 and 2775 of human myeloid/lymphoid or mixed-lineage leukemia 1 using the number given in Swiss-Prot entry Q03164 (GeneID 4297).
Product Application Details	
Applications	Western Blot, Simple Western, Flow Cytometry, Immunoprecipitation, Chromatin Immunoprecipitation (ChIP)
Recommended Dilutions	Western Blot 1:2000-1:10000, Simple Western 1:10, Flow Cytometry, Immunoprecipitation 2-5 ug/mg of lysate, Chromatin Immunoprecipitation (ChIP) 1-5 ug/ml
Application Notes	KMT2A/MLL antibody validated for ELOW from a varified sustamor review

Product Application Details	
Applications	Western Blot, Simple Western, Flow Cytometry, Immunoprecipitation, Chromatin Immunoprecipitation (ChIP)
Recommended Dilutions	Western Blot 1:2000-1:10000, Simple Western 1:10, Flow Cytometry, Immunoprecipitation 2-5 ug/mg of lysate, Chromatin Immunoprecipitation (ChIP) 1-5 ug/ml
Application Notes	KMT2A/MLL antibody validated for FLOW from a verified customer review.




Images


Western Blot: KMT2A/MLL Antibody [NB600-256] - Whole cell lysate (50 ug) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-MLL1 antibody used for WB at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.




Flow Cytometry: KMT2A/MLL Antibody [NB600-256] - Flow cytometry testing of KMT2A/MLL antibody on murine WT and Mll1-/- CD4 T cells that were stimulated in-vitro with anti-CD3/-CD28 antibodies. Image submitted by our verified customer (see Review section for additional details).



Chromatin Immunoprecipitation: KMT2A/MLL Antibody [NB600-256] - A. 10 ug of was used to immunoprecipitate chromatin from K562 cells according to Ren et al (Genes Dev. 2002 16: 245-256).



Immunoprecipitation: KMT2A/MLL Antibody [NB600-256] - Detection of human MLL1 by western blot of immunoprecipitates. Samples: Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HEK293T cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-MLL1 antibody NB600-256 used for IP at 3 ug per reaction. MLL1 was also immunoprecipitated by another rabbit anti-MLL1 antibody. For blotting immunoprecipitated MLL1, NB600-256 was used at 1 ug/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.



Publications

Zhang L, Ye F, Zuo Z et al. Long noncoding RNA TPT1-AS1 promotes the progression and metastasis of colorectal cancer by upregulating the TPT1-mediated FAK and JAK-STAT3 signalling pathways Aging (Albany NY) 2021-02-15 [PMID: 33428595]

Jang M, An J, Oh SW et al. Matrix stiffness epigenetically regulates the oncogenic activation of the Yes-associated protein in gastric cancer Nature biomedical engineering 2020-12-07 [PMID: 33288878]

Schaller M, Ito T, Allen RM et al. Epigenetic regulation of IL-12-dependent T cell proliferation. J Leukoc Biol 2015-06-09 [PMID: 26059830]

Park UH, Yoon SK, Park T et al. Additional sex comb-like (ASXL) proteins 1 and 2 play opposite roles in adipogenesis via reciprocal regulation of peroxisome proliferator-activated receptor {gamma}. J Biol Chem 2011-01-01 [PMID: 21047783]





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NB600-256

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-55237PEP KMT2A/MLL Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB600-256

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

