

Product Datasheet

Ctip1 Antibody - BSA Free NB600-258

Unit Size: 100 ul

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB600-258

Updated 2/21/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB600-258



NB600-258

Ctip1 Antibody - BSA Free

Product Information	
Unit Size	100 ul
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	Tris-Citrate/Phosphate (pH 7.0 - 8.0)
Target Molecular Weight	91 kDa


Product Description	
Host	Rabbit
Gene ID	53335
Gene Symbol	BCL11A
Species	Human
Immunogen	The immunogen recognized by this antibody maps to a region between residues 1 and 50 of human B-cell CLL/Lymphoma 11A using the numbering given in Swiss-Prot entry Q9H165 (GeneID 53335).

Product Application Details	
Applications	Western Blot, Immunoprecipitation
Recommended Dilutions	Western Blot 1:1000-1:10000, Immunoprecipitation 2-10 ug/mg lysate




Images

Western Blot: Ctip1 Antibody [NB600-258] - Detection of Human Ctip1 (Bcl11a) by western blot. Samples: Whole cell lysate (5, 15 and 50 ug) from Ramos cells. Antibodies: Affinity purified rabbit Ctip1 Antibody NB600-258 used at 0.4 ug/ml. Detection: Chemiluminescence with exposure times of 3 minutes.



Immunoprecipitation: Ctip1 Antibody [NB600-258] - Detection of human Ctip1 by western blot of immunoprecipitates. Samples: Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from Ramos cells. Antibodies: Affinity purified rabbit Ctip1 Antibody NB600-258 used for IP at 6 ul/mg lysate. Ctip1 was also immunoprecipitated by rabbit anti-Ctip1 antibodies (NB600-259, NB600-261) and rabbit anti-Ctip1 recombinant monoclonal antibody [BLR073G] (NBP2-78710). For blotting immunoprecipitated Ctip1, NBP2-78710 was used at 1:1000. Chemiluminescence with an exposure time of 75 seconds.



Publications

Chen Z, Luo HY, Steinberg MH, Chui DH. BCL11A represses HBG transcription in K562 cells. Blood Cells Mol Dis 2009-03-01 [PMID: 19153051] (WB, Human)



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NB600-258

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
H00053335-P01-10ug	Recombinant Human Ctip1 GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB600-258

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

