

Product Datasheet

Goat anti-Rat IgG (H+L) Secondary Antibody [FITC] NB7114

Unit Size: 1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB7114

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB7114



NB7114

Goat anti-Rat IgG (H+L) Secondary Antibody [FITC]

Product Information	
Unit Size	1 ml
Concentration	1.0 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.09% Sodium Azide
Isotype	IgG
Conjugate	FITC
Purity	Immunogen affinity purified
Buffer	Phosphate Buffered Saline (PBS) containing 0.2%BSA

Product Description	
Host	Goat
Species	Rat
Specificity/Sensitivity	By immunoelectrophoresis and ELISA this Goat anti-Rat IgG (H+L) Secondary Antibody [FITC] reacts specifically with rat IgG and with light chains common to other rat immunoglobulins. No was detected against non-immunoglobulin serum proteins. This may cross react with IgG from other species.
Immunogen	This Goat anti-Rat IgG (H+L) Secondary Antibody [FITC] was developed against rat IgG-heavy and light chain.

Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Flow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500, Immunohistochemistry-Paraffin 1:50 - 1:500

Publications

Mandakhbayar, N;Ji, Y;El-Fiqi, A;Patel, KD;Yoon, DS;Dashnyam, K;Bayaraa, O;Jin, G;Tsogtbaatar, K;Kim, TH;Lee, JH;Kim, HW; Double hits with bioactive nanozyme based on cobalt-doped nanoglass for acute and diabetic wound therapies through anti-inflammatory and pro-angiogenic functions Bioactive materials 2024-01-01 [PMID: 37637079] (Immunohistochemistry, Immunocytochemistry/ Immunofluorescence)

Laskowski A, Schmidt W, Dinkel K et al. bFGF and EGF modulate trauma-induced proliferation and neurogenesis in juvenile organotypic hippocampal slice cultures. Brain Res. 2005-03-01 [PMID: 15777755]

Details:

This citation used the FITC form of this antibody.

Othman, M et al. Adenovirus-induced thrombocytopenia: the role of von Willebr factor P-selectin in mediating accelerated platelet clearance. Blood. 2007-04-01 [PMID: 17148587]

Details:

This citation used the FITC form of this antibody.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB7114

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

