Product Datasheet

Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] NB7351

Unit Size: 1 ml

Store at 4C. Do not freeze.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NB7351

Updated 10/23/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NB7351



NB7351

Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC]

Product InformationUnit Size1 mlConcentration1.0 mg/mlStorageStore at 4C. Do not freeze.ClonalityPolyclonalPreservative0.09% Sodium AzideConjugateFITCPurityImmunogen affinity purifiedBufferPhosphate Buffered Saline (PBS)Product DescriptionRabbitHostRabbitSpecificity/SensitivityBy immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non- immunoglobulins serum proteins. This may cross react with IgG from other species.ImmunogenThis Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG.Product Application DetailsFlow Cytometry. Immunocytochemistry/ Immunofluorescence, ImmunohistochemistryApplicationsFlow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions": ICC: 1:200-1/2,000 "Optimal dilutions should be determined by the end user.		
Concentration 1.0 mg/ml Storage Store at 4C. Do not freeze. Clonality Polyclonal Preservative 0.09% Sodium Azide Conjugate FITC Purity Immunogen affinity purified Buffer Phosphate Buffered Saline (PBS) Product Description Host Host Rabbit Species Goat Specificity/Sensitivity By immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non-immunoglobulin serum proteins. This may cross react with IgG from other species. Immunogen This Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG. Product Application Details Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry Recommended Dilutions Flow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunohistochemistry Application Notes This antibody can be used in Immunocytochemistry and Immunohistochemistry-Parafin. Recommended Starting Dilutions*': ICC: 1/200-1/2,000 *Optimal	Product Information	
Storage Store at 4C. Do not freeze. Clonality Polyclonal Preservative 0.09% Sodium Azide Conjugate FITC Purity Immunogen affinity purified Buffer Phosphate Buffered Saline (PBS) Product Description Host Host Rabbit Species Goat Specificity/Sensitivity By immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non-immunoglobulin serum proteins. This may cross react with IgG from other species. Immunogen This Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry Recommended Dilutions Flow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry 1:50 - 1:500, Immunocytochemistry and Immunohistochemistry-Paraffin. Recommended Starting Dilutions ¹ : ICC: 1/20.01/2,000 *Optimal	Unit Size	1 ml
ClonalityPolyclonalPreservative0.09% Sodium AzideConjugateFITCPurityImmunogen affinity purifiedBufferPhosphate Buffered Saline (PBS)Product DescriptionHostRabbitSpeciesGoatSpeciesGoatSpecificity/SensitivityBy immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non- immunoglobulin serum proteins. This may cross react with IgG from other 	Concentration	1.0 mg/ml
Preservative 0.09% Sodium Azide Conjugate FITC Purity Immunogen affinity purified Buffer Phosphate Buffered Saline (PBS) Product Description Rabbit Host Rabbit Species Goat Specificity/Sensitivity By immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non- immunoglobulin serum proteins. This may cross react with IgG from other species. Immunogen This Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry Recommended Dilutions Flow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500, Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Storage	Store at 4C. Do not freeze.
Conjugate FITC Purity Immunogen affinity purified Buffer Phosphate Buffered Saline (PBS) Product Description Host Host Rabbit Species Goat Specificity/Sensitivity By immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non- immunoglobulin serum proteins. This may cross react with IgG from other species. Immunogen This Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry Recommended Dilutions Flow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500, Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Clonality	Polyclonal
Purity Immunogen affinity purified Buffer Phosphate Buffered Saline (PBS) Product Description Host Rabbit Species Goat Specificity/Sensitivity By immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non- immunoglobulin serum proteins. This may cross react with IgG from other species. Immunogen This Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry Recommended Dilutions Flow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:200, Immunohistochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Preservative	0.09% Sodium Azide
Buffer Phosphate Buffered Saline (PBS) Product Description Rabbit Host Rabbit Species Goat Specificity/Sensitivity By immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non- immunoglobulin serum proteins. This may cross react with IgG from other species. Immunogen This Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry Recommended Dilutions Flow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500 Application Notes This antibody can be used in Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Conjugate	FITC
Product Description Host Rabbit Species Goat Specificity/Sensitivity By immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non- immunoglobulin serum proteins. This may cross react with IgG from other species. Immunogen This Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG. Product Application Details Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry Recommended Dilutions Flow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Purity	Immunogen affinity purified
HostRabbitSpeciesGoatSpecificity/SensitivityBy immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non- immunoglobulin serum proteins. This may cross react with IgG from other species.ImmunogenThis Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG.Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunohistochemistryRecommended DilutionsFlow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500Application NotesThis antibody can be used in Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Buffer	Phosphate Buffered Saline (PBS)
SpeciesGoatSpecificity/SensitivityBy immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non- immunoglobulin serum proteins. This may cross react with IgG from other species.ImmunogenThis Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG.Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunohistochemistryRecommended DilutionsFlow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500Application NotesThis antibody can be used in Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Product Description	
Specificity/SensitivityBy immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non- immunoglobulin serum proteins. This may cross react with IgG from other species.ImmunogenThis Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG.Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunohistochemistryRecommended DilutionsFlow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500Application NotesThis antibody can be used in Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Host	Rabbit
Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non- immunoglobulin serum proteins. This may cross react with IgG from other species.ImmunogenThis Rabbit anti-Goat IgG (H+L) Secondary Antibody [FITC] was developed against goat IgG.Product Application DetailsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunohistochemistryRecommended DilutionsFlow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500Application NotesThis antibody can be used in Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Species	Goat
against goat IgG.Product Application DetailsApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunohistochemistryRecommended DilutionsFlow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500Application NotesThis antibody can be used in Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Specificity/Sensitivity	Secondary Antibody [FITC] reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No was detected against non-immunoglobulin serum proteins. This may cross react with IgG from other
ApplicationsFlow Cytometry, Immunocytochemistry/ Immunofluorescence, ImmunohistochemistryRecommended DilutionsFlow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500Application NotesThis antibody can be used in Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Immunogen	
ImmunohistochemistryRecommended DilutionsFlow Cytometry 1:50 - 1:200, Immunohistochemistry 1:50 - 1:500, Immunocytochemistry/ Immunofluorescence 1:50 - 1:500Application NotesThis antibody can be used in Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Product Application Details	
Immunocytochemistry/ Immunofluorescence 1:50 - 1:500 Application Notes This antibody can be used in Immunocytochemistry and Immunohistochemistry- Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Applications	
Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal	Recommended Dilutions	
	Application Notes	Paraffin. Recommended Starting Dilutions*: ICC: 1/200-1/2,000 *Optimal





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB7351

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

