

Product Datasheet

Rabbit anti-Goat IgG (H+L) Secondary Antibody [HRP] (Pre-adsorbed) NB7376

Unit Size: 1 ml

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Publications: 1

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NB7376

Updated 10/23/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NB7376



NB7376

Rabbit anti-Goat IgG (H+L) Secondary Antibody [HRP] (Pre-adsorbed)

Product Information	
Unit Size	1 ml
Concentration	0.5 mg/ml
Storage	Store at 4C. Do not freeze.
Clonality	Polyclonal
Preservative	0.05% Pro-Clean 400
Conjugate	HRP
Purity	Immunogen affinity purified
Buffer	Phosphate Buffered Saline (PBS)

Product Description	
Host	Rabbit
Species	Goat
Reactivity Notes	Rat reactivity reported in scientific literature (PMID; 23131471)
Specificity/Sensitivity	By immunoelectrophoresis and ELISA this Rabbit anti-Goat IgG (H+L) Secondary Antibody [HRP] (Pre-adsorbed) reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Antiserum was cross adsorbed using chicken, horse, human, mouse, pig and rat immunosorbents to remove cross reactive Antibodies. Less than 0.1% cross reactivity to chicken, horse, human, mouse, pig, rabbit and rat IgG was detected. This antibody may cross react with IgG from other species.
Immunogen	This Rabbit anti-Goat IgG (H+L) Secondary Antibody [HRP] (Pre-adsorbed) was developed against goat IgG-heavy and light chain.

Product Application Details	
Applications	Western Blot, ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot 1:5000 - 1:50000, ELISA 1:10000- 1:100000, Immunohistochemistry 1:200 - 1:5000, Immunocytochemistry/ Immunofluorescence 1:200 - 1:5000, Immunohistochemistry-Paraffin

Publications

Sun SH, Liu SQ, Cai CP et al. Down-regulation of alpha-2u globulin in renal mitochondria of STZ-induced diabetic rats observed by a proteomic method Ann Endocrinol (Paris) 2012-11-08 [PMID: 23131471]





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Secondary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NB7376

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

