## **Product Datasheet**

### LAS2 Overexpression Lysate NBL1-08256

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-08256

Updated 11/9/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-08256



#### NBL1-08256

LAS2 Overexpression Lysate

Product Information				
Unit Size	0.1 mg			
Concentration	The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.			
Storage	Store at -80C. Avoid freeze-thaw cycles.			
Buffer	RIPA buffer			
Target Molecular Weight	41.6 kDa			
Product Description				
Description	LAS2 Transient Overexpression Lysate			
	Expression Host: HEK293T			
	Plasmid: RC206158			
	Accession#: NM_173529			
	Protein Tag: C-MYC/DDK			
	You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.			
Gene ID	162681			
Gene Symbol	C18ORF54			
Species	Human			
Notes	HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.			
Lysate Type	Overexpression			
Product Application Details				
Applications	Western Blot			
Recommended Dilutions	Western Blot			



Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and
detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

#### innay

Western Blot: C18orf54 Overexpression Lysate (Adult Normal) [NBL1- 08256] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for C18orf54.	158 <del>-</del> 106 - 79 -
	48 —
	35 - 23 -





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBL1-08256

AF5769	MITF Antibody
AF1638	Nucleostemin Antibody [Unconjugated]
NBP2-15638	LAS2 Antibody
H00003845-M01	KRAS Antibody (3B10-2F2)

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-08256

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

