Product Datasheet

COX7C Overexpression Lysate NBL1-09422

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-09422

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-09422



NBL1-09422

0.1 mg
The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.
Store at -80C. Avoid freeze-thaw cycles.
RIPA buffer
5.3 kDa
COX7C Transient Overexpression Lysate
Expression Host: HEK293T
Plasmid: RC201768
Accession#: NM_001867
Protein Tag: C-MYC/DDK
You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.
1350
COX7C
Human

pecies	Hun
--------	-----

Species	пиша
Notes	HEK29

HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid
polymer transfection reagent specially designed and manufactured for large
volume DNA transfection. Transfected cells were cultured for 48hrs before
collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6,
150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM
PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein
concentration was measured by BCA protein assay kit.

Lysate Type	Overexpression
-------------	----------------

Product Application Details Applications Western Blot **Recommended Dilutions** Western Blot




Application Notes

This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (NBP1-71705) present on the protein construct.

Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Images

Western Blot: COX7C Overexpression Lysate (Adult Normal) [NBL1-09422] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for COX7C.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBL1-09422

H00001350-P01-10ug Recombinant Human COX7C GST (N-Term) Protein

NB100-56503 Cytochrome c Antibody (7H8.2C12) - BSA Free

NBP1-98539 COX7C Antibody
DGB300 IGFBP-3 [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-09422

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

