Product Datasheet

FGF basic/FGF2/bFGF Overexpression Lysate NBL1-10694

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-10694

Updated 11/9/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-10694



NBL1-10694

Product Information Unit Size 0.1 mg Concentration The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information Storage Store at -80C. Avoid freeze-thaw cycles. Buffer RIPA buffer Target Molecular Weight 30.6 kDa Product Description Description FGF basic/FGF2/bFGF Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC217426	
Concentration The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information Storage Store at -80C. Avoid freeze-thaw cycles. Buffer RIPA buffer Target Molecular Weight 30.6 kDa Product Description Description FGF basic/FGF2/bFGF Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC217426	
overexpression lysates. Please contact technical support for more information Storage Store at -80C. Avoid freeze-thaw cycles. Buffer RIPA buffer Target Molecular Weight 30.6 kDa Product Description Description FGF basic/FGF2/bFGF Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC217426	
Buffer Target Molecular Weight 30.6 kDa Product Description Description FGF basic/FGF2/bFGF Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC217426	
Target Molecular Weight 30.6 kDa Product Description Description FGF basic/FGF2/bFGF Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC217426	
Product Description Description FGF basic/FGF2/bFGF Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC217426	
FGF basic/FGF2/bFGF Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC217426	
FGF basic/FGF2/bFGF Transient Overexpression Lysate Expression Host: HEK293T Plasmid: RC217426	
Plasmid: RC217426	
A	
Accession#: NM_002006	
Protein Tag: C-MYC/DDK	
You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consist of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue 100mM DTT.	ts
Gene ID 2247	
Gene Symbol FGF2	
Species Human	
Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7. 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1m PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.	
Lysate Type Overexpression	

Dan desart	A I! 1!	on Details
Product	Anniicati	an Detalle
I I Oddot	Applicati	on Details

Applications	Western Blot
Recommended Dilutions	Western Blot




Application Notes

This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag (NBP1-71705) present on the protein construct.

Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Images

Western Blot: FGF-2 Overexpression Lysate (Adult Normal) [NBL1-10694] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for FGF-2.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBL1-10694

NBP2-34921-10ug Recombinant Human FGF basic/FGF2/bFGF Protein

236-EG-200 EGF [Unconjugated]

233-FB-025 FGF basic/FGF2/bFGF [Unconjugated]

210-TA-005 TNF-alpha [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-10694

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

