

Product Datasheet

MRPL21 Overexpression Lysate NBL1-13245

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBL1-13245

Updated 11/9/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBL1-13245



NBL1-13245**MRPL21 Overexpression Lysate**

| Product Information | |
|------------------------------------|--|
| Unit Size | 0.1 mg |
| Concentration | The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information. |
| Storage | Store at -80C. Avoid freeze-thaw cycles. |
| Buffer | RIPA buffer |
| Target Molecular Weight | 22.6 kDa |
| Product Description | |
| Description | <p>MRPL21 Transient Overexpression Lysate</p> <p>Expression Host: HEK293T</p> <p>Plasmid: RC224225</p> <p>Accession#: NM_181514</p> <p>Protein Tag: C-MYC/DDK</p> <p>You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.</p> |
| Gene ID | 219927 |
| Gene Symbol | MRPL21 |
| Species | Human |
| Notes | HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit. |
| Lysate Type | Overexpression |
| Product Application Details | |
| Applications | Western Blot |
| Recommended Dilutions | Western Blot |


Application Notes

This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag ([NBP1-71705](#)) present on the protein construct.

Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

Images

Western Blot: MRPL21 Overexpression Lysate (Adult Normal) [NBL1-13245] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for MRPL21.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBL1-13245

| | |
|---------------|--|
| NBP2-13612PEP | MRPL21 Recombinant Protein Antigen |
| AF2535 | MMR/CD206/Mannose Receptor Antibody [Unconjugated] |
| NBP2-13612 | MRPL21 Antibody - BSA Free |
| NBP2-16756 | Glutathione S-Transferase pi 1/GSTP1 Antibody - BSA Free |

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-13245

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

