## **Product Datasheet**

### PPAR gamma/NR1C3 Overexpression Lysate NBL1-14635

Unit Size: 0.1 mg

Store at -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBL1-14635

Updated 11/9/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBL1-14635



#### NBL1-14635

PPAR gamma/NR1C3 Overexpression Lysate


Unit Size   0.1 mg     Concentration   The exact concentration of the protein of interest cannot be determined for overexpression lysates. Please contact technical support for more information.     Storage   Store at -80C. Avoid freeze-thaw cycles.     Buffer   RIPA buffer     Target Molecular Weight   54.5 kDa     Product Description   Percent State     Description   PPAR gamma/NR1C3 Transient Overexpression Lysate     Expression Host: HEK293T   Plasmid: RC212449     Accession#: NM_005037   Protein Tag: C-MYC/DDK     You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.     Gene ID   5468     Gene Symbol   PPARG     Species   Human     Notes   HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid notwer transfection deal manufactured for large	Product Information	Product Information	
overexpression lysates. Please contact technical support for more information.StorageStore at -80C. Avoid freeze-thaw cycles.BufferRIPA bufferTarget Molecular Weight54.5 kDaProduct DescriptionPPAR gamma/NR1C3 Transient Overexpression LysateDescriptionPPAR gamma/NR1C3 Transient Overexpression LysateView of the state	Unit Size	0.1 mg	
Buffer RIPA buffer   Target Molecular Weight 54.5 kDa   Product Description PPAR gamma/NR1C3 Transient Overexpression Lysate   Expression Host: HEK293T Expression Host: HEK293T   Plasmid: RC212449 Accession#: NM_005037   Protein Tag: C-MYC/DDK You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.   Gene ID 5468   Gene Symbol PPARG   Species Human   Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid	Concentration		
Target Molecular Weight 54.5 kDa   Product Description PPAR gamma/NR1C3 Transient Overexpression Lysate   Description PPAR gamma/NR1C3 Transient Overexpression Lysate   Expression Host: HEK293T Plasmid: RC212449   Accession#: NM_005037 Protein Tag: C-MYC/DDK   You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.   Gene ID 5468   Gene Symbol PPARG   Species Human   Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid	Storage	Store at -80C. Avoid freeze-thaw cycles.	
Product Description   Description   PPAR gamma/NR1C3 Transient Overexpression Lysate   Expression Host: HEK293T   Plasmid: RC212449   Accession#: NM_005037   Protein Tag: C-MYC/DDK   You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.   Gene ID 5468   Gene Symbol PPARG   Species Human   Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid	Buffer	RIPA buffer	
DescriptionPPAR gamma/NR1C3 Transient Overexpression LysateExpression Host: HEK293TPlasmid: RC212449Accession#: NM_005037Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5468Gene SymbolPPARGSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid	Target Molecular Weight	54.5 kDa	
PPAR gamma/NR1C3 Transient Overexpression LysateExpression Host: HEK293TPlasmid: RC212449Accession#: NM_005037Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5468Gene SymbolPPARGSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid	Product Description		
Plasmid: RC212449Accession#: NM_005037Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5468Gene SymbolPPARGSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid	Description		
Accession#: NM_005037Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5468Gene SymbolPPARGSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid		Expression Host: HEK293T	
Protein Tag: C-MYC/DDKYou will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5468Gene SymbolPPARGSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid		Plasmid: RC212449	
You will receive 1 vial of lysate (100ug), 1 vial of empty vector negative control (100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCI pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5468Gene SymbolPPARGSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid		Accession#: NM_005037	
(100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue, 100mM DTT.Gene ID5468Gene SymbolPPARGSpeciesHumanNotesHEK293T cells in 10-cm dishes were transiently transfected with a non-lipid		Protein Tag: C-MYC/DDK	
Gene Symbol PPARG   Species Human   Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid		(100ug), and 1 vial of 2xSDS sample buffer (250ul). Each vial of cell lysate contains 100ug of total protein (at 1 mg/ml). The 2xSDS Sample Buffer consists of 4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromophenol blue,	
Species Human   Notes HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid	Gene ID	5468	
Notes   HEK293T cells in 10-cm dishes were transiently transfected with a non-lipid	Gene Symbol	PPARG	
	Species	Human	
volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCI pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein concentration was measured by BCA protein assay kit.	Notes	polymer transfection reagent specially designed and manufactured for large volume DNA transfection. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4, and then centrifuged to clarify the lysate. Protein	
Lysate Type Overexpression	Lysate Type	Overexpression	
Product Application Details			
Applications Western Blot	Applications	Western Blot	
Recommended Dilutions Western Blot	Recommended Dilutions	Western Blot	



Application Notes	This product is intended for use as a positive control in Western Blot. Overexpression of the target protein was confirmed using an antibody to DDK (FLAG) epitope tag ( <u>NBP1-71705</u> ) present on the protein construct.
	Each vial of cell lysate contains 100ug of total protein which should be sufficient for 20-50 reactions. Depending on over-expression level, antibody affinity and detection system, some lysates can go as low as 0.1 ug per load. We recommend starting with 5ug of cell lysate. Add an equal amount of cell lysate and 2X SDS Sample buffer and boil the SDS samples for 10 minutes before loading.

#### Images

Western Blot: PPAR gamma Overexpression Lysate (Adult Normal) [NBL1-14635] Left-Empty vector transfected control cell lysate (HEK293 cell lysate); Right -Over-expression Lysate for PPAR gamma.



www.novusbio.com





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBL1-14635

NBP2-22106PEP	PPAR gamma/NR1C3 Antibody Blocking Peptide
210-TA-005	TNF-alpha [Unconjugated]
NBP2-22106	PPAR gamma/NR1C3 Antibody - BSA Free
M6000B-1	IL-6 [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Lysates are guaranteed for 6 months from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBL1-14635

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

