

# Product Datasheet

## TGN38 Antibody - Azide and BSA Free NBP1-03495

Unit Size: 0.5 mg

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

**Reviews: 1 Publications: 17**

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP1-03495](http://www.novusbio.com/NBP1-03495)

Updated 2/21/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP1-03495](http://www.novusbio.com/reviews/destination/NBP1-03495)



**NBP1-03495**

TGN38 Antibody - Azide and BSA Free

Product Information	
Unit Size	0.5 mg
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	No Preservative
Reconstitution Instructions	Reconstitute 0.5 mg size in 0.5 ml of sterile water. Centrifuge to remove any insoluble material. Glycerol may be added (1:1) for additional stability. Please note the 0.1 mg size is provided in reconstituted format.
Isotype	IgG
Purity	Ammonium sulfate precipitation
Buffer	Contains 0.02% benzalkonium chloride.


Product Description	
Description	PLEASE NOTE: 0.1mg sample size is provided in reconstituted format and the 0.5mg size is provided lyophilized. (Please see reconstitution instructions).
Host	Rabbit
Gene ID	10618
Gene Symbol	TGOLN2
Species	Human, Mouse, Rat, Monkey
Reactivity Notes	Marmoset (100%).
Marker	TGN Marker
Specificity/Sensitivity	TGN38 (or TGN46, TGN48, TGN51 in human).
Immunogen	A synthetic peptide from mouse TGN38 conjugated to blue carrier protein was used as the antigen.

Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 10-50 ug/ml, Immunohistochemistry 10-50 ug/ml, Immunocytochemistry/ Immunofluorescence 1:50-1:250
Application Notes	The 0.1 mg size ships reconstituted and the 0.5 mg size ships lyophilized. ICC/IF reactivity reported in (PMID: 27505890) and a verified customer review.




## Images


Immunocytochemistry/Immunofluorescence: TGN38 Antibody [NBP1-03495] - Mouse embryonic fibroblasts, fixed with 4% PFA washed with PBS, permeabilized with 0.2% Triton X-100. Primary antibody at 1:100 in 3% BSA in PBS. Secondary antibodies incubation for 2 hours. Image from a verified customer review.




Immunocytochemistry/Immunofluorescence: TGN38 Antibody [NBP1-03495] - Human Melanoma cell line C 32.




Immunocytochemistry/Immunofluorescence: TGN38 Antibody [NBP1-03495] - Human Melanoma cell line C 32



Immunocytochemistry/Immunofluorescence: TGN38 Antibody [NBP1-03495] - Human Melanoma cell line C 32 was cultured overnight on round cover slides placed in a 24 well tissue culture plate. Culture media removed and washed twice with PBS before fixing with 2% formalin for 10 minutes. Cells were then washed three times with PBS and incubated with Tris 0.01M containing Triton X 0.005% for 15 minutes. Cells were washed and incubated with 100 ul of Rabbit antibody to TGN diluted 1:100 in the blocking buffer for 30 minutes. Wells were then washed 7 times with PBS and incubated with 100 ul of anti Rb-FITC conjugate diluted 1:100 in the blocking buffer for further 30 minutes. Cells were washed as before and nuclear counter stained with Hoechst and mounted on to slides.



Immunocytochemistry/Immunofluorescence: TGN38 Antibody [NBP1-03495] - Human Melanoma cell line C 32 was cultured overnight on round cover slides placed in a 24 well tissue culture plate. Culture media removed and washed twice with PBS before fixing with 2% formalin for 10 minutes. Cells were then washed three times with PBS and incubated with Tris 0.01M containing Triton X 0.005% for 15 minutes. Cells were washed and incubated with 100 ul of Rabbit antibody to TGN diluted 1:100 in the blocking buffer for 30 minutes. Wells were then washed 7 times with PBS and incubated with 100 ul of anti Rb-FITC conjugate diluted 1:100 in the blocking buffer for further 30 minutes. Cells were washed as before and nuclear counter stained with Hoechst and mounted on to slides.



## Publications

Saller BS, Wöhrle S, Fischer L et Al. Acute suppression of mitochondrial ATP production prevents apoptosis and provides an essential signal for NLRP3 inflammasome activation Immunity 2024-11-15 [PMID: 39571574]

Tao Yu, Dan Hou, Jiaqi Zhao, Xuan Lu, Wendy K. Greentree, Qian Zhao, Min Yang, Don-Gerard Conde, Maurine E. Linder, Hening Lin NLRP3 Cys126 palmitoylation by ZDHHC7 promotes inflammasome activation Cell reports 2024-05-22 [PMID: 38583156]

Daisuke Sugahara, Yuka Kobayashi, Yoshihiro Akimoto, Hayato Kawakami Mouse intestinal niche cells express a distinct  $\alpha$ 1,2-fucosylated glycan recognized by a lectin from Burkholderia cenocepacia. Glycobiology 2018-06-12 [PMID: 28177462]

Liu Y, Zhai H, Alemayehu H et al. Cryo-electron tomography of NLRP3-activated ASC complexes reveals organelle co-localization Nature communications 2023-11-09 [PMID: 37945612] (ICC/IF, Mouse)

Details:

1:250 ICC/IF dilution

Bilekova S, Garcia-Colomer B, Cebrian-Serrano A et al. Inceptor facilitates acrosomal vesicle formation in spermatids and is required for male fertility Frontiers in Cell and Developmental Biology 2023-08-24 [PMID: 37691832] (Immunohistochemistry-Frozen, Immunocytochemistry/ Immunofluorescence)

Magupalli VG, Negro R, Tian Y et al. HDAC6 mediates an aggresome-like mechanism for NLRP3 and pyrin inflammasome activation Science 2020-09-18 [PMID: 32943500]

Petkovic M, Oses-Prieto JA, Burlingame A, Jan L TMEM16K is an interorganelle regulator of endosomal sorting Nat Commun 2020-07-05 [PMID: 32620747]

Andreeva L, David L, Rawson S et al. NLRP3 cages revealed by full-length mouse NLRP3 structure control pathway activation Cell 2021-12-22 [PMID: 34861190] (ICC/IF, Mouse)

Watson AR, Dai H, Zheng Y et al. mTORC2 Deficiency Alters the Metabolic Profile of Conventional Dendritic Cells Front Immunol 2019-07-02 [PMID: 31338091] (ICC/IF, Mouse)

Shi CS, Nabar NR, Huang NN, Kehrl JH SARS-Coronavirus Open Reading Frame-8b triggers intracellular stress pathways and activates NLRP3 inflammasomes Cell Death Discov 2019-06-05 [PMID: 31231549] (ICC/IF)

Fink, A;Mikulicic, S;Blaum, F;Reddehase, MJ;Florin, L;Lemmermann, NAW; Function of the cargo sorting dileucine motif in a cytomegalovirus immune evasion protein Med. Microbiol. Immunol. 2019-04-19 [PMID: 31004199] (ICC/IF, Mouse)

Ren C, Yuan Q, Braun M et al. Leukocyte Cytoskeleton Polarization Is Initiated by Plasma Membrane Curvature from Cell Attachment Dev. Cell 2019-03-23 [PMID: 30930167] (ICC/IF, Mouse)

More publications at <http://www.novusbio.com/NBP1-03495>



### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

[www.novusbio.com](http://www.novusbio.com)  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP1-03495**

---

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
2914-HT-100MG	Holo-Transferrin [Unconjugated]

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP1-03495](http://www.novusbio.com/reviews/submit/NBP1-03495)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

