

# Product Datasheet

## CTR2 Antibody - BSA Free

### NBP1-05199

Unit Size: 0.1 ml

Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

#### Publications: 4

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP1-05199](http://www.novusbio.com/NBP1-05199)

Updated 9/9/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP1-05199](http://www.novusbio.com/reviews/destination/NBP1-05199)



**NBP1-05199**

CTR2 Antibody - BSA Free

Product Information	
Unit Size	0.1 ml
Concentration	0.93 mg/ml
Storage	Store at 4C short term. Aliquot and store at -20C long term. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.1% Sodium Azide
Isotype	IgG
Purity	Immunogen affinity purified
Buffer	PBS and 30% Glycerol
Target Molecular Weight	71 kDa


Product Description	
Description	Novus Biologicals Rabbit CTR2 Antibody - BSA Free (NBP1-05199) is a polyclonal antibody validated for use in WB and Simple Western. Anti-CTR2 Antibody: Cited in 4 publications. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	1318
Gene Symbol	SLC31A2
Species	Human, Mouse
Immunogen	Synthetic peptide made to an internal portion of human CTR2 (within residues 50-100). [Swiss-Prot# O15432]

Product Application Details	
Applications	Western Blot, Simple Western
Recommended Dilutions	Western Blot 1 ug/ml, Simple Western 10 ug/ml
Application Notes	<p>This CTR2 antibody is useful for Western blot, where a band is seen at approx. 71 kDa. The theoretical molecular weight of CTR2 is 15 kDa. The difference between the expected size and the observed size in Western blot is believed to be due to a multimeric form of CTR2 (Bertinato et al).</p> <p>In Simple Western only 10 - 15 uL of the recommended dilution is used per data point.</p> <p>See <a href="#">Simple Western Antibody Database</a> for Simple Western validation: Tested in MEF and Human Placenta lysate 0.5 mg/mL, separated by Size, antibody dilution of 10 ug/mL. Separated by Size-Wes, Sally Sue/Peggy Sue.</p> <p>The observed molecular weight of the protein may vary from the listed predicted molecular weight due to post translational modifications, post translation cleavages, relative charges, and other experimental factors.</p>




## Images

Western Blot: CTR2 Antibody [NBP1-05199] - Detection of CTR2 in MEF cell lysate.



Simple Western: CTR2 Antibody [NBP1-05199] - Simple Western lane view shows a specific band for CTR2 in 0.5 mg/ml of MEF (left) and Human Placenta (right) lysate. This experiment was performed under reducing conditions using the 12-230 kDa separation system.



## Publications

Solier S, Müller S, Cañeque T et al. A druggable copper-signalling pathway that drives inflammation Nature 2023-04-26 [PMID: 37100912] (WB, Human)

Sun S, Zhao S, Yang Q et al. Enhancer of zeste homolog 2 promotes cisplatin resistance by reducing cellular platinum accumulation Cancer Sci. [PMID: 29630768] (WB, Human)

Blair BG, Larson CA, Adams PL et al. Copper transporter 2 regulates endocytosis and controls tumor growth and sensitivity to cisplatin in vivo. Mol Pharmacol;79(1):157-166. 2011-01-01 [PMID: 20930109] (WB, Mouse)

Blair BG et al. Regulation of CTR2 Expression by Copper and Cisplatin in Human Ovarian Carcinoma Cells. Mol Pharmacol. 2010-03-01 [PMID: 20194531] (WB, Mouse)

## Procedures

### Serum protocol for CTR2 Antibody (NBP1-05199)

CTR2 Antibody:

Western Blot Protocol

1. Perform SDS-PAGE (4-12% MOPS) on samples to be analyzed, loading 20 ug of total protein per lane.
2. Transfer proteins to Nitrocellulose according to the instructions provided by the manufacturer of the transfer apparatus.
3. Rinse membrane with dH<sub>2</sub>O and then stain the blot using Ponceau S for 1-2 minutes to access the transfer of proteins onto the nitrocellulose membrane. Rinse the blot in water to remove excess stain and mark the lane locations and locations of molecular weight markers using a pencil.
4. Rinse the blot in TBS for approximately 5 minutes.
5. Block the membrane using 5% NFD<sub>M</sub> + 1% BSA in TBS + Tween, 1 hour at RT.
6. Rinse the membrane in dH<sub>2</sub>O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
7. Dilute the rabbit anti-CTR2 primary antibody (NBP1-05199) in blocking buffer and incubate 1 hour at room temperature.
8. Rinse the membrane in dH<sub>2</sub>O and then wash the membrane in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each.
9. Apply the diluted rabbit-IgG HRP-conjugated secondary antibody in blocking buffer (as per manufacturers instructions) and incubate 1 hour at room temperature.
10. Wash the blot in wash buffer [TBS + 0.1% Tween] 3 times for 10 minutes each (this step can be repeated as required to reduce background).
11. Apply the detection reagent of choice in accordance with the manufacturers instructions (Pierce ECL).

Note: Tween-20 can be added to the blocking or antibody dilution buffer at a final concentration of 0.05-0.2%, provided it does not interfere with antibody-antigen binding.





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP1-05199**

---

NB800-PC10	MEF Whole Cell Lysate
NBP1-05199PEP	CTR2 Antibody Blocking Peptide
NBP2-33376H	Blue Marker Antibody (6F4-F6) [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP1-05199](http://www.novusbio.com/reviews/submit/NBP1-05199)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)



