

Product Datasheet

LCAT Antibody NBP1-05950

Unit Size: 0.1 ml

Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Reviews: 1 Publications: 3

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP1-05950

Updated 10/23/2024 v.20.1

**Earn rewards for product
reviews and publications.**

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP1-05950




NBP1-05950

LCAT Antibody


Product Information	
Unit Size	0.1 ml
Concentration	This product is unpurified. The exact concentration of antibody is not quantifiable.
Storage	Aliquot and store at -20C or -80C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.05% Sodium Azide
Isotype	IgG
Purity	Unpurified
Buffer	Whole antisera
Product Description	
Host	Rabbit
Gene ID	3931
Gene Symbol	LCAT
Species	Human, Mouse
Immunogen	Carboxyl-terminal histidine-tagged mouse recombinant LCAT purified from the medium of CHO cells overexpressing mouse LCAT cDNA. [UniProt# P16301]
Product Application Details	
Applications	Western Blot, Immunocytochemistry/ Immunofluorescence
Recommended Dilutions	Western Blot 1:1000, Immunocytochemistry/ Immunofluorescence 1:500
Application Notes	In Western blot, a band is seen at ~62-65 kDa representing LCAT. Also has been shown to immunodeplete LCAT from mouse plasma, suggesting it should work fine for Immunoprecipitation (PMID: 18583707).

Images

Western Blot: LCAT Antibody [NBP1-05950] - Analysis of plasma LCAT protein mass. Plasma samples (0.2ul) from four hA-I Tg mice, four hA-I Tg SR-BI ^{-/-} mice, a LCAT^{-/-} mouse, and purified mouse LCAT were fractionated on SDS-PAGE, and LCAT was detected by Western blot analysis using rabbit antiserum against mouse LCAT.



Immunocytochemistry/Immunofluorescence: LCAT Antibody [NBP1-05950] - The LCAT antibody was tested at a 1:500 dilution in HeLa cells against Dylight 488 (Green). Alpha-tubulin and nuclei were counterstained against Dylight 550 (Red) and DAPI (Blue).



Publications

Hirsch-Reinshagen, V et al. LCAT synthesized by primary astrocytes esterifies cholesterol on glia-derived lipoproteins. *J Lipid Res*;50(5):885-93. 2009-05-01 [PMID: 19065001]

Mulya, A et al. Initial interaction of apoA-I with ABCA1 impacts in vivo metabolic fate of nascent HDL. *J Lipid Res*;49(11):2390-401. 2008-11-01 [PMID: 18583707]

Lee, J et al. Functional LCAT deficiency in human apolipoprotein A-I transgenic, SR-BI knockout mice. *J Lipid Res*;48(5):1052-61. 2007-05-01 [PMID: 17272829]



Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP1-05950

NBL1-12457	LCAT Overexpression Lysate
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP1-05950

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

